

**SOCIO-ECONOMIC DIFFERENCES
IN
HEALTH, NUTRITION, AND POPULATION**

**MALI
1995/96, 2001**

**Davidson R. Gwatkin, Shea Rutstein, Kiersten Johnson,
Eldaw Suliman, Adam Wagstaff, and Agbessi Amouzou**

TABLE OF CONTENTS

Foreword	I
Introduction	III
Part I. Basic Tables, 2001	1
A. Total Population	3
B. Female and Male Populations	9
C. Rural and Urban Populations	13
Part II. Basic Tables, 1995 / 96	19
A. Total Population	21
B. Female and Male Populations	27
C. Rural and Urban Populations	31
Part III. Technical Notes	37
A. Indicator Definitions	39
B. Data and Methods	53
C. Discussion	57
Part IV. Supporting Tables, 2001	63
A. Sample Sizes	65
B. Standard Errors	71
C. Asset Distribution and Weights	77
Part V. Supporting Tables, 1995 / 96	79
A. Sample Sizes	81
B. Standard Errors	87
C. Asset Distribution and Weights	93

Part VI. Annexes	95
A. Sources of Additional Information	97
B. Use of Information from this Report to Monitor the Economic Status of People Served by HNP Programs	99
C. Countries Covered by the HNP-Poverty Report Project	105

FOREWORD

The World Bank shares the desire of its member states and client countries to ensure that the poor partake fully in the health gains that the countries achieve. To assist in this, the Bank, in cooperation with the Dutch and Swedish Governments, has sponsored the set of reports providing basic information about health inequalities within countries to which this document belongs.

The information shows clearly that disparities in both health conditions and health service use are unacceptably large. As countries and the Bank work to reduce important inequalities among regions and countries, there is a clear need for equally vigorous efforts to lessen the inequity represented by intra-country differences among socio-economic groups.

My colleagues and I hope that all concerned with equity in health will find this information useful in making the case for effective actions to improve the health of the poor, and in designing programs to achieve this crucial objective.

Joy Phumaphi
Vice President
Human Development Network
The World Bank

INTRODUCTION

This report is one in a series that provides basic information about health, nutrition, and population (hnp) inequalities within fifty-six developing countries.

The series to which the report belongs is an expanded and updated version of a set covering forty-five countries that was published in 2000. The fifty-six reports in the current series cover almost all DHS surveys undertaken during the period beginning in 1990 and ending with the date of the last survey for which data were publicly available as of June 2006.¹

The report's contents are intended to facilitate preparation of country analyses and the development of activities to benefit poor people. To this end, the report presents data about hnp status, service use, and related matters among individuals belonging to different socio-economic classes. The principal focus is on differences among groups of individuals defined in terms of the wealth or assets of the households where they reside. The source of data is the Demographic and Health Survey (DHS) program, a large, multi-country household survey project.

The figures in this and the other reports in the series draw on responses to questions about household wealth or assets included in the DHS questionnaire, which were similar for all the surveys covered. These responses served as the basis for the construction of a wealth index, which was used to rank individuals according to the index value for the household to which they belonged. The individuals were then divided into quintiles, and the mean value for each of up to approximately 120 indicators was calculated for each quintile.

The report is organized in four principal parts:

- Parts one and two, which constitute the report's core, consist of tables presenting quintile-specific data for each hnp indicator covered from the two most recent DHS surveys available at the time of the report's publication. In each part there are three sets of tables: the first provides quintile-specific information for the total population; the second presents data separately for quintiles of females and males; the third features quintile-specific information presented separately for rural and urban residents. Each of these sets is divided into four sections: one dealing with hnp status, the second with the use of hnp services, the third with hnp-related behavior, and the fourth with other hnp status determinants.

- Part three provides supplementary technical information designed to help readers understand the data presented in parts one and two. This information deals with such issues as how the covered hnp indicators were defined and how the quintile-specific estimates were derived.

- Parts four and five present supporting tables that deal with three of the technical matters covered in part three: the size of the sample for each indicator covered; the standard error

¹ The average interval of approximately two years between data collection and availability means that the latest surveys covered were conducted in 2005.

for each quintile-specific estimate in the total population; and the items used in constructing the wealth index, along with the weight assigned to each.

An additional, sixth part consists of three annexes, for readers interested in applying the approach used in the report or in learning more about the other reports in this series. The first annex is an annotated bibliography containing further information about the technical issues concerning the approach used in the report, and about employing that approach to examine additional issues using DHS or other data sets. The second shows how the report's approach can be applied to monitor the distribution of benefits from other hnp programs, and provides a tool for doing so. The third annex is a list of all fifty-six countries for which reports are available, along with an indication of how to obtain copies of the reports dealing with them.

The authors thank the Dutch and Swedish Governments for the generous support that made production of this report possible.

PART I. BASIC TABLES, 2001

A. TOTAL POPULATION

B. FEMALE AND MALE POPULATIONS

C. RURAL AND URBAN POPULATIONS

Notes:

- Each of the three sections referred to above consists of four divisions, presenting data for: I) hnp status; II) hnp service use; III) hnp-related individual and household behavior; and IV) other, underlying determinants of hnp status.
- Full definitions of all indicators covered in the tables are provided in section A of the technical notes found in part II.
- “na” appears in the table cells when data are not available, usually because the DHS survey concerned did not collect information about the indicator(s) in question.
- Figures in the tables shown within parentheses indicate the absence of adequate observations to produce acceptably reliable values. Asterisks appear when the number of observations was too small to justify the presentation even of figures within parentheses. (For further information, see the section on “Sampling Errors” in the presentation of data and methods in part II.B.) Asterisks also will be found in columns showing statistical indices of inequality when the amount of quintile-specific information available is inadequate to permit computation of the value for the index concerned.
- Female/male tables include only indicators relevant for both sexes; those pertaining to only one sex (e.g., fertility, women’s nutritional status, antenatal care, attended deliveries) have been omitted.

Mali
2001 - TOTAL POPULATION

Part I: HNP STATUS

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Childhood illness and mortality										
Infant mortality rate	137.2	125.2	140.6	128.7	89.9	126.2	1.53	47.30	-0.0492	0.0006
Under-five mortality rate	247.8	264.1	262.0	244.4	148.1	238.2	1.67	99.70	-0.0618	0.0008
Prevalence of fever	29.5	26.9	28.1	25.0	24.1	26.8	1.22	5.40	-0.0385	0.0096
Prevalence of diarrhea	20.8	20.5	20.7	16.9	13.0	18.6	1.60	7.80	-0.0725	0.0117
Prevalence of acute respiratory infection (ARI)	10.4	9.0	10.8	8.4	10.9	9.9	0.95	0.50	0.0436	0.0184
B. Fertility										
Total fertility rate	7.3	7.5	7.3	6.8	5.3	6.8	1.38	2.00	-0.0591	0.0003
Adolescent fertility rate	200.0	246.0	225.0	203.0	108.0	185.0	1.85	92.00	-0.1295	0.0016
C. Nutritional status (%)										
<i>Children:</i>										
Moderate stunting	19.7	22.4	19.0	18.4	12.2	18.5	1.61	7.50	-0.0888	0.0122
Severe stunting	25.1	23.8	21.4	15.7	7.5	19.1	3.35	17.60	-0.1681	0.0117
Moderate underweight	26.0	27.3	24.4	20.4	12.8	22.5	2.03	13.20	-0.1139	0.0108
Severe underweight	13.2	14.1	12.3	9.3	3.9	10.8	3.38	9.30	-0.1529	0.0164
Mild anemia	16.1	19.0	15.7	20.1	21.6	18.2	0.75	5.50	0.0329	0.0248
Moderate anemia	56.7	52.2	57.7	51.2	44.5	53.0	1.27	12.20	-0.0344	0.0111
Severe anemia	14.1	11.2	13.8	11.9	4.3	11.5	3.28	9.80	-0.1357	0.0320
<i>Women:</i>										
Malnutrition	13.0	13.7	13.5	13.3	10.2	12.6	1.27	2.80	-0.0579	0.0149
Mild anemia	42.8	41.0	38.1	40.3	33.3	39.1	1.29	9.50	-0.0248	0.0120
Moderate anemia	21.9	24.6	25.2	20.7	13.4	20.8	1.63	8.50	-0.1044	0.0180
Severe anemia	2.7	4.2	2.8	2.4	2.2	2.8	1.23	0.50	-0.1195	0.0589
D. Female circumcision (%)										
<i>Prevalence of circumcision:</i>										
Girls	76.3	73.3	66.0	67.9	79.2	72.4	0.96	2.90	-0.0253	0.0039
Women	95.9	93.8	89.7	88.4	90.6	91.6	1.06	5.30	-0.0202	0.0016
<i>Prevalence of occlusion:</i>										
Girls	3.4	3.7	5.8	4.7	5.2	4.5	0.65	1.80	-0.0279	0.0374
Women	1.3	1.4	2.0	1.9	2.6	1.9	0.50	1.30	0.0836	0.0420
E. Sexually transmitted disease										
<i>Prevalence of genital discharge:</i>										
Women	20.4	17.8	18.6	20.1	29.9	21.5	0.68	9.50	0.0806	0.0110
Men	4.3	5.5	5.5	4.3	4.1	4.7	1.05	0.20	-0.0285	0.0476
<i>Prevalence of genital ulcer:</i>										
Women	8.2	9.0	8.0	9.1	9.9	8.8	0.83	1.70	0.0158	0.0177
Men	2.9	3.6	1.1	1.7	1.3	2.0	2.23	1.60	-0.1419	0.0801

Mali

2001 - TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Childhood immunization										
BCG coverage	58.1	60.4	59.6	77.6	92.1	69.0	0.63	34.00	0.0995	0.0080
Measles coverage	39.7	39.3	38.3	52.8	76.5	48.7	0.52	36.80	0.1396	0.0133
DPT coverage	28.1	28.7	28.6	45.7	70.7	39.6	0.40	42.60	0.2096	0.0156
Full basic coverage	19.5	20.2	18.7	31.9	56.0	28.7	0.35	36.50	0.2250	0.0210
No basic coverage	29.4	25.8	27.7	17.6	6.1	21.7	4.82	23.30	-0.1924	0.0226
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	na	na	na	na	na	na	na	na	na	na
B. Treatment of childhood illnesses										
<i>Treatment of fever:</i>										
Medical treatment of fever	18.4	18.9	25.4	29.2	61.1	28.9	0.30	42.70	0.2649	0.0167
Treatment in a public facility	6.5	7.1	11.4	14.2	43.4	15.1	0.15	36.90	0.4135	0.0284
Treatment in a private facility	11.7	11.8	13.6	14.8	17.3	13.6	0.68	5.60	0.0983	0.0272
<i>Treatment of acute respiratory infection (ARI):</i>										
Medical treatment of ARI	21.6	28.9	26.7	33.6	69.7	35.6	0.31	48.10	0.2510	0.0232
Treatment in a public facility	8.9	9.6	13.0	22.3	49.6	20.3	0.18	40.70	0.4127	0.0378
Treatment in a private facility	12.5	19.3	12.8	11.3	19.3	14.9	0.65	6.80	0.0329	0.0439
<i>Treatment of diarrhea:</i>										
Use of oral rehydration therapy	64.4	63.7	60.6	66.9	79.8	65.7	0.81	15.40	0.0144	0.0094
Medical treatment of diarrhea	12.3	11.7	12.2	16.6	42.5	16.8	0.29	30.20	0.2120	0.0310
Treatment in a public facility	2.9	2.8	4.3	9.3	27.5	7.5	0.11	24.60	0.4276	0.0536
Treatment in a private facility	9.3	8.8	7.9	7.4	15.0	9.2	0.62	5.70	0.0303	0.0399
C. Antenatal and delivery care										
<i>Antenatal care visits:</i>										
To a medically trained person	42.2	44.0	46.2	64.6	91.8	56.8	0.46	49.60	0.1676	0.0052
To a doctor	0.9	1.4	1.8	2.5	7.8	2.7	0.12	6.90	0.4477	0.0471
To a nurse or trained midwife	41.4	42.6	44.4	62.0	84.0	54.1	0.49	42.60	0.1521	0.0057
Multiple visits to a medically trained person	24.7	29.3	30.8	45.8	73.8	40.0	0.33	49.10	0.2294	0.0076
<i>Antenatal care content:</i>										
Tetanus toxoid	35.1	37.4	37.7	55.0	81.3	48.5	0.43	46.20	0.1725	0.0064
Prophylactic antimalarial treatment	48.5	51.1	52.7	65.7	86.6	60.2	0.56	38.10	0.1174	0.0051
Iron supplementation	25.5	25.5	27.1	40.2	57.2	34.5	0.45	31.70	0.1420	0.0124
<i>Delivery attendance:</i>										
By a medically trained person	22.0	27.0	29.4	45.8	88.7	40.6	0.25	66.70	0.2769	0.0059
By a doctor	0.4	0.6	1.1	1.2	4.8	1.5	0.08	4.40	0.4471	0.0538
By a nurse or trained midwife	21.5	26.4	28.3	44.6	83.9	39.1	0.26	62.40	0.2704	0.0062
In a public facility	19.7	24.1	26.4	42.7	80.1	36.8	0.25	60.40	0.2821	0.0067
In a private facility	0.4	0.5	0.2	0.5	3.8	1.0	0.11	3.40	0.4801	0.0752
At home	78.9	73.7	72.1	55.7	14.4	60.9	5.48	64.50	-0.1769	0.0040
D. Contraceptive services										
<i>Contraceptive prevalence:</i>										
Women	4.2	3.6	3.4	7.3	17.9	7.0	0.23	13.70	0.3423	0.0250
Men	7.6	5.6	6.2	12.7	19.1	10.2	0.40	11.50	0.2941	0.0431

Mali

2001 - TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
D. Contraceptive services (cont.)										
<i>Source of contraception - public sector:</i>										
Women	33.3	29.9	49.2	54.8	49.1	46.3	0.68	15.80	0.0500	0.0228
Men	na	na	na	na	na	na	na	na	na	na
<i>Source of contraception - private sector:</i>										
Women	19.5	32.9	34.9	31.1	44.9	36.6	0.43	25.40	0.1507	0.0258
Men	na	na	na	na	na	na	na	na	na	na
E. Treatment of adult illnesses										
<i>Treatment of genital discharge, ulcer, sore:</i>										
Women	14.7	19.0	15.5	31.6	60.2	32.4	0.24	45.50	0.3067	0.0148
Men	(19.7)	(20.0)	(17.7)	(32.7)	(34.7)	25.7	0.57	15.00	0.2305	0.0771
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>										
Women	0.8	1.0	1.5	3.6	12.0	4.2	0.07	11.20	0.5528	0.0308
Men	4.6	3.7	3.8	7.0	19.2	8.7	0.24	14.60	0.3983	0.0398

Mali

2001 - TOTAL POPULATION

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Hygienic practices										
<i>Disposal of children's stools:</i>										
Sanitary disposal	na	na	na	na	na	na	na	na	na	na
<i>Handwashing:</i>										
Wash hands prior to preparing food	90.6	90.1	88.3	89.8	92.7	90.4	0.98	2.10	0.0058	0.0017
Handwashing facilities in household	3.0	4.6	7.3	7.6	19.5	8.2	0.15	16.50	0.3604	0.0195
B. Bednet ownership and use										
<i>Bednet ownership:</i>										
Bednet ownership	52.7	56.8	50.8	54.1	58.8	54.4	0.90	6.10	0.0335	0.0049
Treated bednet ownership	na	na	na	na	na	na	na	na	na	na
<i>Bednet use:</i>										
By children	34.4	38.6	34.9	37.1	48.0	38.3	0.72	13.60	0.0748	0.0086
By pregnant women	33.4	36.6	34.6	36.7	39.4	35.9	0.85	6.00	0.0738	0.0200
C. Breastfeeding										
Exclusive breastfeeding	38.0	32.2	26.5	24.0	17.9	28.2	2.12	20.10	-0.1152	0.0296
Timely complementary feeding	30.1	24.9	27.7	35.8	44.3	32.4	0.68	14.20	0.0703	0.0294
Bottle-feeding	2.0	2.8	1.9	3.2	12.4	4.2	0.16	10.40	0.4041	0.0710
D. Micronutrient consumption										
<i>Iodized salt:</i>										
Availability of iodized salt in household	40.2	35.2	29.9	33.7	51.9	37.2	0.77	11.70	0.0018	0.0089
<i>Vitamin A:</i>										
Children	34.3	34.6	39.3	42.0	56.2	40.9	0.61	21.90	0.1107	0.0077
Women	9.5	13.3	13.4	22.2	32.1	17.7	0.30	22.60	0.2331	0.0143
E. Tobacco and alcohol use										
<i>Tobacco:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
<i>Alcohol:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
F. Sexual practices										
<i>Non-regular sexual partnerships:</i>										
Women	0.4	0.3	0.5	0.7	1.9	0.8	0.21	1.50	0.3978	0.0679
Men	2.4	4.4	6.2	8.5	12.1	7.1	0.20	9.70	0.2745	0.0387
<i>Condom usage with non-regular partner:</i>										
Women	*	*	*	*	34.1	22.3	*	*	*	*
Men	36.3	33.8	33.9	70.7	65.1	55.5	0.56	28.80	0.1542	0.0293
G. Domestic violence										
Ever experienced violence	na	na	na	na	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na	na	na	na	na

Mali

2001 - TOTAL POPULATION

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Education										
<i>School completion:</i>										
Women	2.4	2.8	3.4	10.0	41.8	13.2	0.06	39.40	0.6143	0.0151
Men	8.3	8.3	10.6	20.3	58.3	24.6	0.14	50.00	0.4779	0.0094
<i>School participation:</i>										
Girls	17.9	18.0	19.5	29.2	65.6	28.3	0.27	47.70	0.3014	0.0130
Boys	28.8	24.2	29.1	39.8	74.1	37.4	0.39	45.30	0.2093	0.0105
B. Exposure to mass media										
<i>Newspaper readership:</i>										
Women	0.4	0.5	0.9	3.1	20.0	5.6	0.02	19.60	0.7002	0.0279
Men	1.6	3.2	4.6	7.8	38.2	13.3	0.04	36.60	0.6377	0.0325
<i>Radio listenership:</i>										
Women	49.7	54.1	51.1	64.8	82.7	61.5	0.60	33.00	0.1099	0.0040
Men	76.3	76.5	80.7	85.6	90.7	82.6	0.84	14.40	0.0429	0.0047
<i>Television viewership:</i>										
Women	11.2	11.6	12.1	29.6	79.1	31.0	0.14	67.90	0.4408	0.0077
Men	22.2	22.7	23.4	43.5	85.6	43.3	0.26	63.40	0.3164	0.0106
C. Knowledge and attitudes about HIV/AIDS										
<i>Knowledge about sexual transmission of HIV/AIDS:</i>										
Women	42.7	41.4	39.3	52.3	80.5	52.6	0.53	37.83	0.1485	0.0048
Men	71.8	69.7	68.8	77.4	88.3	76.3	0.81	16.50	0.0607	0.0055
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>										
Women	32.6	33.5	31.8	43.6	74.5	45.2	0.44	41.90	0.1687	0.0058
Men	52.9	53.5	52.2	62.8	76.2	62.4	0.69	23.30	0.0658	0.0078
<i>Attitudes toward HIV/AIDS:</i>										
Women	48.2	48.9	47.5	43.9	29.5	43.1	1.63	18.70	-0.1084	0.0062
Men	55.1	56.4	53.8	46.0	27.7	46.3	1.99	27.40	-0.1408	0.0104
D. Status of women										
<i>Household decisionmaking:</i>										
Can seek own health care	9.7	8.9	12.4	12.3	13.6	11.5	0.71	3.90	0.1008	0.0151
Can seek children's health care	63.4	64.7	65.7	77.5	83.9	70.7	0.76	20.50	0.0696	0.0043
Can make daily household purchases	26.8	23.8	27.1	25.1	23.7	25.2	1.13	3.10	-0.0231	0.0092
Can make large household purchases	15.1	13.2	18.5	17.3	18.0	16.5	0.84	2.90	0.0455	0.0122
Can make meal-related decisions	50.8	46.9	49.4	44.1	40.2	46.0	1.26	10.60	-0.0517	0.0058
<i>Freedom of movement:</i>										
Can travel to visit family, relatives	15.8	14.6	18.9	22.1	23.5	19.2	0.67	7.70	0.0957	0.0114
<i>Other decisionmaking, attitudes:</i>										
Can decide how to spend own money	82.6	82.5	84.4	84.3	86.6	84.4	0.95	4.00	0.0124	0.0033
Can decide whether to have sex	72.7	75.4	73.5	73.1	75.5	74.1	0.96	2.80	0.0102	0.0031
Justifies domestic violence	88.9	90.9	90.2	90.6	84.2	88.8	1.06	4.70	-0.0086	0.0021
E. Orphanhood										
Paternal orphan prevalence	2.5	3.0	4.1	4.1	5.3	3.7	0.47	2.80	0.1339	0.0182
Maternal orphan prevalence	2.0	2.2	2.1	1.8	2.4	2.1	0.83	0.40	0.0537	0.0237
Double orphan prevalence	0.4	0.7	0.6	0.5	0.6	0.6	0.67	0.20	0.1231	0.0464

Mali
2001 - FEMALE / MALE POPULATIONS

Part I: HNP STATUS

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood mortality and morbidity												
Infant mortality rate	131.3	113.3	126.3	119.3	77.2	115.6	143.4	136.7	154.3	137.4	102.0	136.4
Under-five mortality rate	233.7	258.1	243.6	233.0	135.3	226.0	262.3	269.8	279.6	255.0	160.3	250.1
Prevalence of fever	31.4	27.1	27.6	23.1	22.6	26.6	27.5	26.7	28.6	26.8	25.5	27.0
Prevalence of diarrhea	19.5	20.0	21.3	14.3	13.1	17.8	22.2	21.1	20.0	19.3	13.0	19.3
Prevalence of acute respiratory infection (ARI)	11.5	9.6	11.2	9.0	10.0	10.3	9.3	8.3	10.5	7.9	11.7	9.5
B. Nutritional status												
<i>Children:</i>												
Moderate stunting	18.9	23.7	18.4	18.1	10.6	18.2	20.6	21.2	19.6	18.7	13.7	18.9
Severe stunting	23.4	24.0	22.1	14.6	7.5	18.8	27.0	23.5	20.8	16.8	7.4	19.4
Moderate underweight	24.8	26.4	23.9	18.3	11.0	21.2	27.3	28.2	25.0	22.4	14.6	23.7
Severe underweight	12.4	14.4	12.5	10.1	4.2	11.0	14.1	13.8	12.1	8.7	3.7	10.7
Mild anemia	16.6	21.8	17.4	17.8	23.2	19.1	15.6	16.4	13.8	21.7	20.1	17.4
Moderate anemia	55.1	51.9	60.8	48.6	44.2	52.8	58.4	52.5	54.3	53.2	44.8	53.2
Severe anemia	12.7	12.1	11.9	15.2	6.2	11.8	15.6	10.4	15.8	9.5	2.4	11.2
C. Sexually transmitted disease												
<i>Prevalence of genital discharge:</i>												
Women	20.4	17.8	18.6	20.1	29.9	21.5						
Men							4.3	5.5	5.5	4.3	4.1	4.7
<i>Prevalence of genital ulcer:</i>												
Women	8.2	9.0	8.0	9.1	9.9	8.8						
Men							2.9	3.6	1.1	1.7	1.3	2.0

Mali

2001 - FEMALE / MALE POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization												
BCG coverage	55.0	59.1	63.4	77.3	92.9	68.7	61.3	61.7	55.8	78.0	91.3	69.3
Measles coverage	36.9	36.7	41.1	51.3	78.3	48.0	42.7	41.6	35.4	54.2	74.8	49.4
DPT coverage	29.3	24.7	27.5	40.8	72.6	38.0	26.7	32.2	29.8	50.2	68.9	41.2
Full basic coverage	22.0	17.0	18.8	32.0	60.0	29.1	17.0	23.2	18.6	31.9	52.4	28.3
No basic coverage	31.5	25.3	27.4	18.6	5.5	22.2	27.3	26.3	28.0	16.8	6.8	21.2
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	na	na	na	na	na	na	na	na	na	na	na	na
B. Treatment of childhood illnesses												
<i>Treatment of fever:</i>												
Medical treatment of fever	17.2	20.7	25.4	32.4	57.7	28.4	19.8	17.0	25.4	26.6	64.0	29.4
Treatment in a public facility	6.8	7.0	11.1	16.0	40.1	14.4	6.0	7.2	11.7	12.8	46.2	15.7
Treatment in a private facility	10.2	13.7	13.7	16.4	17.2	13.8	13.5	9.9	13.5	13.5	17.3	13.4
<i>Treatment of acute respiratory infection (ARI):</i>												
Medical treatment of ARI	18.5	34.5	24.4	31.4	65.9	33.2	25.5	22.5	29.2	35.9	72.8	38.2
Treatment in a public facility	8.2	12.4	13.0	20.3	46.7	18.8	9.8	6.3	13.1	24.4	52.0	21.9
Treatment in a private facility	10.0	22.0	9.9	11.1	18.5	13.9	15.7	16.2	15.7	11.5	19.9	16.0
<i>Treatment of diarrhea:</i>												
Use of oral rehydration therapy	66.3	62.5	64.2	67.5	83.4	67.2	62.7	64.9	56.8	66.6	76.2	64.4
Medical treatment of diarrhea	10.3	14.8	10.8	13.0	42.5	16.2	14.2	8.8	13.8	19.1	42.5	17.4
Treatment in a public facility	2.2	2.3	4.0	4.2	28.2	6.5	3.5	3.2	4.7	12.7	26.8	8.4
Treatment in a private facility	8.0	12.2	6.8	8.9	14.3	9.6	10.4	5.6	9.0	6.4	15.7	8.9
C. Contraceptive services												
<i>Contraceptive prevalence:</i>												
Women	4.2	3.6	3.4	7.3	17.9	7.0						
Men							7.6	5.6	6.2	12.7	19.1	10.2
<i>Source of contraception - public sector:</i>												
Women	33.3	29.9	49.2	54.8	49.1	46.3						
Men							na	na	na	na	na	na
<i>Source of contraception - private sector:</i>												
Women	19.5	32.9	34.9	31.1	44.9	36.6						
Men							na	na	na	na	na	na
D. Treatment of adult illnesses												
<i>Treatment of genital discharge, ulcer, sore:</i>												
Women	14.7	19.0	15.5	31.6	60.2	32.4						
Men							(19.7)	(20.0)	(17.7)	(32.7)	(34.7)	0.0
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>												
Women	0.8	1.0	1.5	3.6	12.0	4.2						
Men							4.6	3.7	3.8	7.0	19.2	8.7

Mali

2001 - FEMALE / MALE POPULATIONS

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Hygienic practices												
<i>Disposal of children's stools:</i>												
Sanitary disposal	na	na	na	na	na	na						
<i>Handwashing:</i>												
Wash hands prior to preparing food	90.6	90.1	88.3	89.8	92.7	90.4						
Handwashing facilities in household	3.0	4.6	7.3	7.6	19.5	8.2						
B. Bednet ownership and use												
<i>Bednet use:</i>												
By children	34.4	38.6	34.9	37.1	48.0	38.3						
C. Breastfeeding												
Exclusive breastfeeding	32.7	21.7	27.7	20.0	18.2	24.5	43.8	42.9	25.1	28.6	17.8	31.9
Timely complementary feeding	31.1	25.8	25.3	28.4	42.8	31.0	28.7	24.1	29.7	41.1	46.2	33.8
Bottle-feeding	3.2	2.9	1.8	4.7	10.2	4.5	0.6	2.7	1.9	1.7	14.7	4.0
D. Micronutrient consumption												
<i>Vitamin A:</i>												
Children	35.0	34.3	39.8	42.4	54.1	40.7	33.6	34.8	38.9	41.7	58.3	41.0
E. Tobacco and alcohol use												
<i>Tobacco:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
<i>Alcohol:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
F. Sexual practices												
<i>Non-regular sexual partnerships:</i>												
Women	0.4	0.3	0.5	0.7	1.9	0.8						
Men							2.4	4.4	6.2	8.5	12.1	7.1
<i>Condom usage with non-regular partner:</i>												
Women	*	*	*	*	34.1	22.3						
Men							36.3	33.8	33.9	70.7	65.1	55.5

Mali

2001 - FEMALE / MALE POPULATIONS

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Education												
<i>School completion:</i>												
Women	2.4	2.8	3.4	10.0	41.8	13.2	8.3	8.3	10.6	20.3	58.3	24.6
Men												
<i>School participation:</i>												
Girls	17.9	18.0	19.5	29.2	65.6	28.3						
Boys							28.8	24.2	29.1	39.8	74.1	37.4
B. Exposure to mass media												
<i>Newspaper readership:</i>												
Women	0.4	0.5	0.9	3.1	20.0	5.6						
Men							1.6	3.2	4.6	7.8	38.2	13.3
<i>Radio listenership:</i>												
Women	49.7	54.1	51.1	64.8	82.7	61.5						
Men							76.3	76.5	80.7	85.6	90.7	82.6
<i>Television viewership:</i>												
Women	11.2	11.6	12.1	29.6	79.1	31.0						
Men							22.2	22.7	23.4	43.5	85.6	43.3
C. Knowledge and attitudes about HIV/AIDS												
<i>Knowledge about sexual transmission of HIV/AIDS:</i>												
Women	42.7	41.4	39.3	52.3	80.5	52.6						
Men							71.8	69.7	68.8	77.4	88.3	76.3
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>												
Women	32.6	33.5	31.8	43.6	74.5	45.2						
Men							52.9	53.5	52.2	62.8	76.2	62.4
<i>Attitudes toward HIV/AIDS:</i>												
Women	48.2	48.9	47.5	43.9	29.5	43.1						
Men							55.1	56.4	53.8	46.0	27.7	46.3
D. Orphanhood												
Paternal orphan prevalence	2.8	3.1	4.0	4.3	5.1	3.8	2.3	2.9	4.2	3.8	5.5	3.7
Maternal orphan prevalence	2.2	2.2	2.4	1.7	2.9	2.3	1.9	2.2	1.9	2.0	1.9	2.0
Double orphan prevalence	0.4	0.7	0.7	0.5	0.7	0.6	0.4	0.7	0.4	0.5	0.6	0.5

Mali
2001 - RURAL / URBAN POPULATIONS

Part I: HNP STATUS

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood illness and mortality												
Infant mortality rate	137.8	125.3	140.6	130.7	66.2	131.9	*	*	(141.5)	123.4	93.5	105.9
Under-five mortality rate	246.2	262.2	263.0	254.4	(138.9)	253.2	*	*	(252.3)	217.3	149.4	184.6
Prevalence of fever	29.4	26.3	28.3	26.6	25.1	27.7	(32.7)	38.0	25.4	20.7	23.9	24.0
Prevalence of diarrhea	20.8	20.5	21.4	18.2	17.7	20.3	(23.2)	20.9	11.8	13.4	12.3	13.1
Prevalence of acute respiratory infection	10.5	8.7	11.0	9.5	13.4	10.1	(8.0)	14.2	8.3	5.4	10.5	9.3
B. Fertility												
Total fertility rate	7.3	7.6	7.3	7.1	*	7.3	*	*	*	6.2	5.2	5.5
Adolescent fertility rate	197.0	250.0	229.0	206.0	*	218.0	*	*	*	195.0	106.0	129.0
C. Nutritional status												
<i>Children:</i>												
Moderate stunting	19.4	21.8	19.7	19.7	13.0	20.0	(33.8)	35.2	10.6	14.9	12.1	13.9
Severe stunting	25.2	24.4	21.6	17.1	11.1	22.1	(19.1)	10.5	19.5	11.8	6.9	9.2
Moderate underweight	25.9	27.4	24.5	21.6	12.3	24.7	(33.3)	26.3	23.2	17.3	12.9	15.5
Severe underweight	13.2	14.4	12.5	10.2	7.0	12.6	(14.2)	8.5	9.4	7.0	3.5	5.1
Mild anemia	16.4	19.3	15.3	18.5	17.5	17.3	*	(12.9)	(20.3)	25.3	22.2	21.7
Moderate anemia	56.5	51.4	57.8	52.2	53.2	54.6	*	(69.0)	(56.0)	48.1	43.3	47.0
Severe anemia	13.7	11.6	13.1	13.3	3.9	12.7	*	(4.2)	(21.9)	7.3	4.4	6.9
<i>Women:</i>												
Malnutrition	12.9	13.6	13.4	12.9	11.2	13.1	(18.9)	14.9	14.3	14.0	10.1	11.4
Mild anemia	42.8	41.6	39.0	40.3	27.7	40.8	*	(30.6)	(25.9)	40.2	33.9	35.0
Moderate anemia	21.8	24.6	24.9	21.2	14.7	22.8	*	(24.6)	(29.7)	19.4	13.2	15.9
Severe anemia	2.8	4.4	2.4	2.6	1.6	3.0	*	0.0	(7.4)	1.8	2.2	2.2
D. Female circumcision												
<i>Prevalence of circumcision:</i>												
Girls	76.2	73.1	66.0	67.7	67.1	71.0	(80.9)	76.1	65.8	68.3	80.9	76.7
Women	95.8	94.2	90.4	88.7	92.0	92.5	100.0	86.8	81.7	87.8	90.4	89.5
<i>Prevalence of occlusion:</i>												
Girls	3.4	3.8	5.8	4.5	3.0	4.3	(4.7)	3.1	5.5	5.1	5.5	5.3
Women	1.2	1.5	2.0	2.5	0.7	1.7	5.6	0.0	1.9	0.5	2.8	2.3
E. Sexually transmitted disease												
<i>Prevalence of genital discharge:</i>												
Women	19.9	17.9	18.6	18.2	22.2	18.8	37.5	16.0	17.6	24.9	30.9	28.4
Men	4.3	5.4	5.2	4.3	4.1	4.7	*	(7.1)	(8.6)	4.5	4.1	4.6
<i>Prevalence of genital ulcer:</i>												
Women	8.3	9.1	8.2	8.5	6.1	8.5	2.8	7.5	5.7	10.4	10.3	9.8
Men	2.8	3.1	1.0	1.8	2.9	2.3	*	(10.4)	(2.1)	1.4	1.1	1.5

Mali

2001 - RURAL / URBAN POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization												
BCG coverage	57.4	59.4	58.2	70.9	95.5	62.0	*	(77.6)	(79.6)	93.3	91.5	90.4
Measles coverage	39.5	39.3	38.1	46.7	67.1	41.3	*	(39.7)	(40.6)	67.0	78.0	70.8
DPT coverage	27.4	27.9	27.1	35.8	59.3	30.1	*	(42.3)	(52.0)	68.5	72.5	68.4
Full basic coverage	18.8	19.7	18.8	27.2	47.1	21.5	*	(30.3)	(17.7)	43.1	57.5	50.3
No basic coverage	29.9	26.5	29.2	23.0	0.0	26.5	*	(14.0)	(5.8)	5.1	7.1	7.1
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	na	na	na	na	na	na	na	na	na	na	na	na
B. Treatment of childhood diseases												
<i>Treatment of fever:</i>												
Medical treatment of fever	18.0	17.5	23.3	24.5	48.8	21.3	*	(39.0)	56.1	46.0	63.1	56.9
Treatment in a public facility	6.4	6.6	11.5	11.8	21.9	9.2	*	(14.1)	11.4	23.0	46.9	36.8
Treatment in a private facility	11.4	10.8	11.7	12.5	27.0	12.0	*	(24.9)	41.3	23.0	15.7	19.6
<i>Treatment of acute respiratory infection (ARI):</i>												
Medical treatment of ARI	20.7	27.0	25.0	31.3	(44.2)	26.2	*	*	(56.7)	(44.8)	74.7	68.4
Treatment in a public facility	8.8	9.5	13.0	20.4	(22.0)	12.8	*	*	(14.2)	(31.8)	55.1	46.4
Treatment in a private facility	11.7	17.5	11.6	10.9	(22.2)	13.3	*	*	(32.2)	(13.0)	18.7	20.7
<i>Treatment of diarrhea:</i>												
Use of oral rehydration therapy	63.9	63.3	60.6	64.8	71.6	63.3	*	*	(60.7)	75.0	81.6	78.0
Medical treatment of diarrhea	11.4	11.4	11.2	12.5	39.1	12.3	*	*	(37.6)	32.4	43.3	39.4
Treatment in a public facility	2.9	2.7	3.7	5.3	15.7	3.8	*	*	(19.9)	24.3	30.1	25.9
Treatment in a private facility	8.4	8.5	7.5	7.2	23.4	8.4	*	*	(17.8)	8.0	13.1	13.3
C. Antenatal and delivery care												
<i>Antenatal care visits:</i>												
To a medically trained person	41.1	42.8	45.0	57.5	92.1	47.2	(86.7)	67.6	61.7	83.4	91.8	86.9
To a doctor	0.8	1.4	1.4	2.2	8.1	1.6	(2.2)	1.7	7.1	3.6	7.8	6.4
To a nurse or trained midwife	40.2	41.4	43.6	55.4	84.0	45.6	(84.5)	65.9	54.6	79.8	84.0	80.5
Multiple visits to a medically trained person	24.1	28.1	29.3	40.6	65.0	30.8	(48.4)	52.9	49.2	59.6	75.0	68.5
<i>Antenatal care content:</i>												
Tetanus toxoid	34.1	36.2	36.2	47.6	78.0	39.1	(72.9)	60.0	57.6	74.8	81.8	77.7
Prophylactic antimalarial treatment	48.2	50.3	51.8	63.4	83.5	53.6	(62.2)	65.5	64.1	71.8	87.1	80.8
Iron supplementation	24.8	24.9	26.2	36.3	55.9	28.3	(54.1)	36.3	39.3	50.6	57.4	53.8
<i>Delivery attendance:</i>												
By a medically trained person	7.2	7.2	9.0	15.1	48.8	10.4	43.0	44.5	43.2	67.1	86.7	76.5
By a doctor	0.4	0.6	1.0	0.7	5.1	0.8	3.4	1.1	2.6	2.6	4.8	3.9
By a nurse or trained midwife	6.9	6.6	8.0	14.4	43.8	9.6	39.7	43.4	40.6	64.6	82.0	72.6
In a public facility	18.4	22.7	24.1	33.2	56.6	24.9	72.9	53.8	55.1	69.3	83.6	76.9
In a private facility	0.4	0.5	0.2	0.4	5.3	0.5	0.0	0.3	0.0	0.8	3.6	2.5
At home	80.2	75.1	74.4	65.5	34.9	73.3	25.8	46.0	43.3	28.5	11.4	19.3
D. Contraceptive services												
<i>Contraceptive prevalence:</i>												
Women	4.3	3.5	3.1	5.4	13.9	4.3	1.7	4.0	6.8	12.4	18.5	15.3
Men	7.7	4.9	6.2	10.0	18.4	7.4	*	*	(7.0)	19.0	19.2	18.2

Mali

2001 - RURAL / URBAN POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
D. Contraceptive services (cont.)												
<i>Source of contraception - public sector:</i>												
Women	32.7	27.2	48.9	55.3	(47.4)	41.3	*	*	*	54.1	49.3	50.6
Men	(100.0)	*	*	(100.0)	*	100.0	0.0	*	*	*	100.0	100.0
<i>Source of contraception - private sector:</i>												
Women	19.7	33.2	32.3	22.6	(38.5)	27.4	*	*	*	40.9	45.5	44.5
Men	na	na	na	na	na	na	na	na	na	na	na	na
E. Treatment of adult illnesses												
<i>Treatment of genital discharge, ulcer, sore:</i>												
Women	14.4	17.8	14.7	27.1	60.6	19.5	*	(41.3)	(25.4)	39.8	60.1	54.1
Men	(20.1)	(23.6)	(19.6)	(22.9)	*	22.1	*	*	*	*	(33.3)	32.9
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>												
Women	0.8	0.8	1.3	1.7	9.9	1.4	0.0	5.0	3.8	8.3	12.2	10.6
Men	4.6	2.5	3.1	2.7	11.6	3.7	*	(21.1)	12.5	15.7	20.1	18.7

Mali

2001 - RURAL / URBAN POPULATIONS

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Hygienic practices												
<i>Disposal of children's stools:</i>												
Sanitary disposal	na	na	na	na	na	na	na	na	na	na	na	na
<i>Handwashing:</i>												
Wash hands prior to preparing food	90.5	90.3	88.2	89.3	89.3	89.6	97.2	87.2	88.8	90.9	93.0	92.3
Handwashing facilities in household	3.0	4.8	7.6	7.1	21.8	6.1	0.0	1.1	4.0	8.9	19.2	14.7
B. Bednet ownership and use												
<i>Bednet ownership:</i>												
Bednet ownership	52.3	56.3	50.0	53.4	68.7	53.4	(70.0)	66.5	59.8	55.7	57.4	57.6
Treated bednet ownership	na	na	na	na	na	na	na	na	na	na	na	na
<i>Bednet use:</i>												
By children	34.1	38.0	33.6	35.5	54.3	35.8	(51.4)	48.2	53.2	41.4	47.0	46.2
By pregnant women	33.2	36.2	34.9	34.7	(57.8)	35.2	*	*	*	43.6	37.3	38.6
C. Breastfeeding												
Exclusive breastfeeding	38.3	32.3	26.4	23.9	*	30.6	*	*	*	(24.5)	16.6	19.5
Timely complementary feeding	30.1	25.0	26.5	35.4	(47.8)	29.7	0.0	*	*	(36.9)	43.6	40.9
Bottle-feeding	2.0	2.5	2.0	2.8	12.2	2.6	*	*	0.0	4.5	12.5	9.7
D. Micronutrient consumption												
<i>Iodized salt:</i>												
Availability of iodized salt in household	40.1	35.6	29.5	30.8	41.8	34.1	(48.3)	27.8	33.9	42.5	53.4	47.6
<i>Vitamin A:</i>												
Children	33.7	33.1	37.6	38.8	51.3	36.1	(60.7)	62.5	61.2	50.6	57.0	56.1
Women	9.2	12.5	12.5	17.4	32.3	13.1	(21.4)	30.5	24.2	34.9	32.1	31.9
E. Tobacco and alcohol use												
<i>Tobacco:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
<i>Alcohol:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
F. Sexual practices												
<i>Non-regular sexual partnerships:</i>												
Women	0.4	0.3	0.5	0.6	0.3	0.4	0.0	0.0	0.0	1.2	2.0	1.7
Men	2.4	4.0	5.1	7.2	20.3	5.0	*	(10.6)	19.4	11.2	11.3	11.5
<i>Condom usage with non-regular partner:</i>												
Women	*	*	*	*	*	(1.3)	*	*	*	*	34.6	35.0
Men	37.8	26.1	36.2	72.9	63.7	47.3	*	(75.9)	26.3	68.0	65.4	62.6
G. Domestic violence												
Ever experienced violence	na	na	na	na	na	na	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na	na	na	na	na	na	na

Mali

2001 - RURAL / URBAN POPULATIONS

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Education												
<i>School completion:</i>												
Women	2.4	2.6	3.0	6.6	27.3	4.3	2.5	6.7	7.7	18.4	43.4	34.7
Men	8.1	7.9	10.0	15.1	47.7	11.4	(22.0)	15.1	18.2	31.3	59.4	50.9
<i>School participation:</i>												
Girls	18.0	17.7	19.4	27.2	54.1	21.0	*	23.7	20.0	34.6	67.3	53.5
Boys	29.0	23.6	28.6	34.2	56.2	29.3	*	(40.1)	35.9	53.4	77.6	66.3
B. Exposure to mass media												
<i>Newspaper readership:</i>												
Women	0.4	0.4	0.9	2.3	18.3	1.5	0.0	2.1	1.2	5.0	20.1	15.3
Men	1.6	3.0	4.6	7.3	17.8	4.2	*	(6.0)	5.4	9.0	40.5	31.7
<i>Radio listenership:</i>												
Women	49.5	52.7	50.5	60.4	81.4	53.8	57.1	76.4	57.9	75.2	82.8	79.4
Men	76.5	75.7	80.9	85.0	92.1	79.4	*	(88.1)	78.4	86.8	90.5	89.0
<i>Television viewership:</i>												
Women	11.3	10.8	11.2	22.4	60.7	15.0	7.1	23.4	23.1	46.7	81.1	68.2
Men	22.2	20.3	22.3	32.3	75.5	25.6	*	(56.7)	36.4	65.7	86.7	79.0
C. Knowledge and attitudes about HIV/AIDS												
<i>Knowledge about sexual transmission of HIV/AIDS:</i>												
Women	46.5	45.9	44.7	50.3	76.7	47.7	36.9	56.3	46.2	69.5	82.6	76.5
Men	72.0	69.5	68.8	74.9	77.6	71.4	*	(71.5)	69.9	82.3	89.5	86.3
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>												
Women	35.6	31.9	31.1	39.0	72.4	35.5	30.0	59.4	39.7	54.6	74.7	67.8
Men	59.5	52.6	51.8	57.2	66.8	56.1	*	(66.6)	57.2	73.9	77.2	75.1
<i>Attitudes toward HIV/AIDS:</i>												
Women	50.8	49.2	47.3	45.0	36.0	47.8	50.8	45.7	49.3	41.4	28.8	33.1
Men	55.2	56.5	53.2	49.9	14.4	52.6	*	(55.4)	(61.6)	38.4	29.1	33.8
D. Status of women												
<i>Household decisionmaking:</i>												
Can seek own health care	9.9	9.0	11.6	12.0	7.3	10.5	2.8	6.3	21.5	13.2	14.3	14.0
Can seek children's health care	62.7	64.3	64.8	74.5	75.4	66.4	(90.7)	71.1	78.6	86.1	85.1	84.4
Can make daily household purchases	27.1	24.1	26.6	25.1	27.3	25.8	15.0	18.6	33.5	25.1	23.3	23.8
Can make large household purchases	15.2	13.2	17.6	17.3	18.2	15.8	11.7	14.0	28.8	17.2	18.0	18.2
Can make meal-related decisions	50.8	47.1	48.5	45.3	47.7	48.1	50.8	43.1	60.3	41.1	39.4	41.1
<i>Freedom of movement:</i>												
Can travel to visit family, relatives	16.0	14.5	18.4	22.4	17.9	17.6	11.4	15.9	24.1	21.4	24.1	23.1
<i>Other decisionmaking, attitudes:</i>												
Can decide how to spend own money	82.3	82.3	85.0	82.9	79.6	83.0	(96.9)	(86.4)	78.8	87.3	87.3	87.0
Can decide whether to have sex	73.0	74.8	73.2	71.8	66.7	73.0	62.1	83.9	77.0	76.3	76.4	76.5
Justifies domestic violence	88.6	91.0	90.5	90.9	87.5	90.1	100.0	90.0	87.4	89.7	83.9	85.7
E. Orphanhood												
Paternal orphan prevalence	2.5	3.0	4.1	3.6	5.2	3.3	3.2	2.0	4.0	5.2	5.3	5.0
Maternal orphan prevalence	2.1	2.2	2.0	1.9	0.7	2.0	0.7	2.7	4.0	1.7	2.7	2.5
Double orphan prevalence	0.4	0.7	0.5	0.4	0.5	0.5	0.0	0.7	1.6	0.9	0.7	0.8

PART II. BASIC TABLES, 1995 / 96

A. TOTAL POPULATION

B. FEMALE AND MALE POPULATIONS

C. RURAL AND URBAN POPULATIONS

Notes:

- Each of the three sections referred to above consists of four divisions, presenting data for: I) hnp status; II) hnp service use; III) hnp-related individual and household behavior; and IV) other, underlying determinants of hnp status.
- Full definitions of all indicators covered in the tables are provided in section A of the technical notes found in part II.
- “na” appears in the table cells when data are not available, usually because the DHS survey concerned did not collect information about the indicator(s) in question.
- Figures in the tables shown within parentheses indicate the absence of adequate observations to produce acceptably reliable values. Asterisks appear when the number of observations was too small to justify the presentation even of figures within parentheses. (For further information, see the section on “Sampling Errors” in the presentation of data and methods in part II.B.) Asterisks also will be found in columns showing statistical indices of inequality when the amount of quintile-specific information available is inadequate to permit computation of the value for the index concerned.
- Female/male tables include only indicators relevant for both sexes; those pertaining to only one sex (e.g., fertility, women’s nutritional status, antenatal care, attended deliveries) have been omitted.

Mali
1995 / 96 - TOTAL POPULATION

Part I: HNP STATUS

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Childhood illness and mortality										
Infant mortality rate	151.4	146.9	138.9	129.0	93.2	133.5	1.62	58.20	-0.0753	0.0006
Under-five mortality rate	297.9	283.6	251.8	241.1	169.1	252.2	1.76	128.80	-0.0901	0.0006
Prevalence of fever	41.9	41.4	40.2	38.8	30.3	38.7	1.38	11.60	-0.0462	0.0104
Prevalence of diarrhea	29.0	26.8	27.8	25.7	16.2	25.3	1.79	12.80	-0.0788	0.0135
Prevalence of acute respiratory infection (ARI)	15.4	16.6	15.0	15.3	13.9	15.3	1.11	1.50	-0.0183	0.0191
B. Fertility										
Total fertility rate	6.9	7.3	7.4	7.1	5.1	6.7	1.35	1.80	-0.0512	0.0005
Adolescent fertility rate	198.0	219.0	210.0	217.0	122.0	187.0	1.62	76.00	-0.0874	0.0017
C. Nutritional status (%)										
<i>Children:</i>										
Moderate stunting	14.2	17.5	16.6	17.0	12.1	15.6	1.17	2.10	-0.0270	0.0189
Severe stunting	18.0	17.7	15.1	13.8	7.2	14.5	2.50	10.80	-0.1360	0.0197
Moderate underweight	25.9	27.4	23.0	21.3	19.6	23.5	1.32	6.30	-0.0618	0.0151
Severe underweight	21.3	18.3	16.5	16.8	8.5	16.5	2.51	12.80	-0.1322	0.0185
Mild anemia	na	na	na	na	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na	na	na	na	na
<i>Women:</i>										
Malnutrition	15.9	18.3	16.1	18.1	12.2	16.2	1.30	3.70	-0.0482	0.0206
Mild anemia	na	na	na	na	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na	na	na	na	na
D. Female circumcision (%)										
<i>Prevalence of circumcision:</i>										
Girls	67.6	74.1	76.5	78.9	79.1	75.2	0.85	11.50	0.0221	0.0048
Women	88.0	95.7	96.5	95.8	92.8	93.7	0.95	4.80	0.0005	0.0026
<i>Prevalence of occlusion:</i>										
Girls	na	na	na	na	na	na	na	na	na	na
Women	0.2	1.0	0.4	0.4	0.5	0.5	0.40	0.30	0.0201	0.0774
E. Sexually transmitted disease										
<i>Prevalence of genital discharge:</i>										
Women	0.1	0.4	0.2	0.4	0.2	0.3	0.50	0.10	0.0541	0.1068
Men	2.1	1.5	2.9	2.3	2.3	2.2	0.91	0.20	0.1239	0.0876
<i>Prevalence of genital ulcer:</i>										
Women	0.2	0.2	0.2	0.1	0.2	0.2	1.00	0.00	0.0545	0.1347
Men	1.9	3.1	0.6	3.2	0.8	1.9	2.38	1.10	-0.1052	0.0820

Mali

1995 / 96 - TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Childhood immunization										
BCG coverage	56.5	70.7	76.2	83.5	94.7	76.3	0.60	38.20	0.0908	0.0075
Measles coverage	34.2	41.3	48.8	52.3	79.7	50.8	0.43	45.50	0.1564	0.0135
DPT coverage	17.4	31.2	31.2	42.4	67.3	37.5	0.26	49.90	0.2481	0.0173
Full basic coverage	15.8	26.2	26.9	34.6	55.6	31.5	0.28	39.80	0.2380	0.0203
No basic coverage	42.5	29.3	22.8	14.5	4.9	22.8	8.67	37.60	-0.3048	0.0248
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	0.5	1.3	3.1	8.0	27.1	7.6	0.02	26.60	0.6136	0.0604
B. Treatment of childhood illnesses										
<i>Treatment of fever:</i>										
Medical treatment of fever	7.6	8.3	14.7	15.3	36.0	14.9	0.21	28.34	0.3317	0.0336
Treatment in a public facility	4.7	4.9	6.8	10.1	18.5	8.3	0.25	13.81	0.2839	0.0482
Treatment in a private facility	2.9	3.4	7.9	5.2	17.5	6.6	0.17	14.53	0.3914	0.0542
<i>Treatment of acute respiratory infection (ARI):</i>										
Medical treatment of ARI	18.2	19.3	28.2	35.7	68.2	32.3	0.27	49.95	0.2738	0.0292
Treatment in a public facility	13.0	11.2	12.4	23.4	43.5	19.6	0.30	30.54	0.2452	0.0441
Treatment in a private facility	5.3	8.1	15.8	12.3	24.7	12.7	0.21	19.41	0.3178	0.0542
<i>Treatment of diarrhea:</i>										
Use of oral rehydration therapy	38.4	43.6	38.6	48.6	67.0	45.0	0.57	28.60	0.0912	0.0170
Medical treatment of diarrhea	8.5	19.3	17.6	21.6	35.1	18.8	0.24	26.68	0.2249	0.0333
Treatment in a public facility	4.2	11.9	10.2	13.3	19.9	11.0	0.21	15.72	0.2306	0.0473
Treatment in a private facility	4.3	7.4	7.4	8.3	15.2	7.8	0.28	10.95	0.2165	0.0567
C. Antenatal and delivery care										
<i>Antenatal care visits:</i>										
To a medically trained person	20.6	35.2	44.2	58.0	85.7	47.5	0.24	65.10	0.2349	0.0078
To a doctor	0.1	0.6	0.7	1.4	2.9	1.1	0.03	2.80	0.5373	0.1014
To a nurse or trained midwife	20.4	34.6	43.5	56.6	82.7	46.4	0.25	62.30	0.2457	0.0075
Multiple visits to a medically trained person	10.8	21.3	31.7	43.8	69.6	34.3	0.16	58.80	0.3209	0.0098
<i>Antenatal care content:</i>										
Tetanus toxoid	31.2	41.2	47.5	59.8	82.1	51.4	0.38	50.90	0.1863	0.0071
Prophylactic antimalarial treatment	na	na	na	na	na	na	na	na	na	na
Iron supplementation	na	na	na	na	na	na	na	na	na	na
<i>Delivery attendance:</i>										
By a medically trained person	11.1	22.8	34.1	51.5	80.6	38.7	0.14	69.50	0.3428	0.0080
By a doctor	0.1	0.3	0.3	1.2	2.0	0.7	0.05	1.90	0.4922	0.1112
By a nurse or trained midwife	11.0	22.6	33.9	50.3	78.6	38.0	0.14	67.60	0.3401	0.0082
In a public facility	8.4	16.0	23.6	40.0	67.6	29.9	0.12	59.20	0.3761	0.0107
In a private facility	0.1	0.0	0.2	0.2	2.3	0.5	0.04	2.20	0.7097	0.1537
At home	89.6	78.6	67.2	51.2	19.9	62.7	4.50	69.70	-0.2074	0.0049
D. Contraceptive services										
<i>Contraceptive prevalence:</i>										
Women	0.5	1.4	2.0	4.2	15.3	4.5	0.03	14.80	0.6124	0.0381
Men	1.0	3.9	6.5	14.3	18.7	8.4	0.05	17.70	0.4168	0.0486

Mali

1995 / 96 - TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
D. Contraceptive services (cont.)										
<i>Source of contraception - public sector:</i>										
Women	*	*	(32.6)	53.9	61.8	57.2	*	*	*	*
Men	na	na	na	na	na	na	na	na	na	na
<i>Source of contraception - private sector:</i>										
Women	*	*	(67.4)	39.0	35.6	39.3	*	*	*	*
Men	na	na	na	na	na	na	na	na	na	na
E. Treatment of adult illnesses										
<i>Treatment of genital discharge, ulcer, sore:</i>										
Women	*	*	*	*	(65.8)	46.4	*	*	*	*
Men	*	*	*	*	*	31.5	*	*	*	*
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na

Mali

1995 / 96 - TOTAL POPULATION

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Hygienic practices										
<i>Disposal of children's stools:</i>										
Sanitary disposal	na	na	na	na	na	na	na	na	na	na
<i>Handwashing:</i>										
Wash hands prior to preparing food	na	na	na	na	na	na	na	na	na	na
Handwashing facilities in household	na	na	na	na	na	na	na	na	na	na
B. Bednet ownership and use										
<i>Bednet ownership:</i>										
Bednet ownership	na	na	na	na	na	na	na	na	na	na
Treated bednet ownership	na	na	na	na	na	na	na	na	na	na
<i>Bednet use:</i>										
By children	na	na	na	na	na	na	na	na	na	na
By pregnant women	na	na	na	na	na	na	na	na	na	na
C. Breastfeeding										
Exclusive breastfeeding	10.9	13.5	12.8	13.7	10.4	12.3	1.05	0.50	-0.0576	0.0603
Timely complementary feeding	23.6	27.9	36.8	32.9	45.8	33.0	0.52	22.20	0.1181	0.0330
Bottle-feeding	1.5	1.1	1.8	4.0	8.8	3.2	0.17	7.30	0.4913	0.0932
D. Micronutrient consumption										
<i>Iodized salt:</i>										
Availability of iodized salt in household	na	na	na	na	na	na	na	na	na	na
<i>Vitamin A:</i>										
Children	1.7	0.5	0.8	1.1	1.0	1.0	1.70	0.70	0.0561	0.1041
Women	na	na	na	na	na	na	na	na	na	na
E. Tobacco and alcohol use										
<i>Tobacco:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
<i>Alcohol:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
F. Sexual practices										
<i>Non-regular sexual partnerships:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	8.1	10.7	16.1	20.7	39.3	19.9	0.21	31.20	0.3203	0.0246
<i>Condom usage with non-regular partner:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	6.2	13.7	18.3	32.2	52.2	35.5	0.12	46.00	0.2865	0.0317
G. Domestic violence										
Ever experienced violence	na	na	na	na	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na	na	na	na	na

Mali

1995 / 96 - TOTAL POPULATION

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles						Low/High Ratio	Low-High Diff. (Abs. Val.)	Concentration Index	
	Low	2nd	3rd	4th	High	Avg.			Value	Standard Errors
A. Education										
<i>School completion:</i>										
Women	1.4	2.4	3.3	8.7	36.1	10.9	0.04	34.70	0.6303	0.0201
Men	4.6	6.9	9.7	19.3	52.6	20.6	0.09	48.00	0.5111	0.0125
<i>School participation:</i>										
Girls	5.0	10.6	14.8	26.9	55.2	22.0	0.09	50.20	0.4590	0.0166
Boys	11.8	14.9	21.7	33.2	62.9	28.0	0.19	51.10	0.3478	0.0140
B. Exposure to mass media										
<i>Newspaper readership:</i>										
Women	0.5	0.9	1.3	4.9	21.7	6.5	0.02	21.20	0.6886	0.0298
Men	3.1	4.5	6.6	14.0	44.1	15.8	0.07	41.00	0.5493	0.0306
<i>Radio listenership:</i>										
Women	29.1	50.6	59.0	70.2	85.3	59.8	0.34	56.20	0.1799	0.0045
Men	59.0	76.8	83.1	90.3	93.9	81.3	0.63	34.90	0.0853	0.0059
<i>Television viewership:</i>										
Women	6.2	12.9	17.3	30.9	73.2	29.8	0.08	67.00	0.4124	0.0089
Men	9.1	15.7	29.5	44.5	82.7	38.3	0.11	73.60	0.3617	0.0129
C. Knowledge and attitudes about HIV/AIDS										
<i>Knowledge about sexual transmission of HIV/AIDS:</i>										
Women	63.7	77.9	79.2	85.0	93.7	80.4	0.68	30.00	0.0661	0.0029
Men	90.7	94.4	95.2	94.8	99.0	95.0	0.92	8.30	0.0156	0.0028
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>										
Women	29.6	26.7	32.6	47.2	73.5	40.8	0.40	43.90	0.2724	0.0064
Men	46.6	43.5	50.0	60.1	77.3	55.7	0.60	30.70	0.1427	0.0102
<i>Attitudes toward HIV/AIDS:</i>										
Women	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na
D. Status of women										
<i>Household decisionmaking:</i>										
Can seek own health care	na	na	na	na	na	na	na	na	na	na
Can seek children's health care	na	na	na	na	na	na	na	na	na	na
Can make daily household purchases	na	na	na	na	na	na	na	na	na	na
Can make large household purchases	na	na	na	na	na	na	na	na	na	na
Can make meal-related decisions	na	na	na	na	na	na	na	na	na	na
<i>Freedom of movement:</i>										
Can travel to visit family, relatives	na	na	na	na	na	na	na	na	na	na
<i>Other decisionmaking, attitudes:</i>										
Can decide how to spend own money	69.9	77.5	78.6	82.8	87.5	80.5	0.80	17.62	0.0364	0.0047
Can decide whether to have sex	na	na	na	na	na	na	na	na	na	na
Justifies domestic violence	na	na	na	na	na	na	na	na	na	na
E. Orphanhood										
Paternal orphan prevalence	3.8	3.6	3.4	4.1	4.4	3.9	0.86	0.60	0.0523	0.0217
Maternal orphan prevalence	2.5	1.7	2.4	2.0	2.1	2.1	1.19	0.40	0.0046	0.0278
Double orphan prevalence	0.4	0.3	0.4	0.3	0.7	0.4	0.57	0.30	0.1167	0.0698

Mali
1995 / 96 - FEMALE / MALE POPULATIONS

Part I: HNP STATUS

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood mortality and morbidity												
Infant mortality rate	150.1	141.8	131.4	115.6	84.7	126.5	152.9	152.2	146.3	142.1	101.6	140.5
Under-five mortality rate	299.7	274.5	243.3	238.2	161.5	247.1	296.3	292.7	260.1	243.6	176.7	257.3
Prevalence of fever	39.5	43.2	37.8	38.5	26.5	37.4	44.6	39.7	42.6	39.1	34.2	40.1
Prevalence of diarrhea	27.3	27.0	26.4	27.2	13.5	24.6	30.9	26.7	29.2	24.3	19.0	26.1
Prevalence of acute respiratory infection (ARI)	16.1	16.8	14.4	15.4	12.7	15.1	14.6	16.4	15.6	15.3	15.0	15.4
B. Nutritional status												
<i>Children:</i>												
Moderate stunting	14.2	18.5	14.4	16.9	11.9	15.3	14.2	16.5	18.9	17.0	12.3	15.9
Severe stunting	16.7	16.2	14.4	14.1	7.1	13.9	19.5	19.3	15.8	13.5	7.3	15.1
Moderate underweight	25.9	28.7	22.2	20.1	22.3	23.9	25.8	26.1	23.8	22.5	17.2	23.1
Severe underweight	21.4	18.0	14.5	17.7	8.7	16.4	21.2	18.6	18.6	16.0	8.4	16.6
Mild anemia	na	na	na	na	na	na	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na	na	na	na	na	na	na
C. Sexually transmitted disease												
<i>Prevalence of genital discharge:</i>												
Women	0.1	0.4	0.2	0.4	0.2	0.3						
Men							2.1	1.5	2.9	2.3	2.3	2.2
<i>Prevalence of genital ulcer:</i>												
Women	0.2	0.2	0.2	0.1	0.2	0.2						
Men							1.9	3.1	0.6	3.2	0.8	1.9

Mali

1995 / 96 - FEMALE / MALE POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization												
BCG coverage	53.0	67.6	73.8	85.0	93.6	74.4	60.7	73.5	79.2	81.8	95.8	78.3
Measles coverage	29.9	36.1	46.8	52.2	72.4	47.0	39.4	46.0	51.3	52.4	87.1	54.9
DPT coverage	16.1	29.1	30.0	44.7	68.8	37.0	19.0	33.1	32.7	39.9	65.8	38.2
Full basic coverage	14.6	24.4	24.3	36.2	57.1	30.7	17.3	27.8	30.1	32.9	54.1	32.4
No basic coverage	46.4	32.4	25.0	13.7	6.4	25.0	37.8	26.5	20.0	15.3	3.4	20.5
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	0.0	0.9	3.1	8.2	25.5	7.0	1.1	1.7	3.0	7.8	28.7	8.2
B. Treatment of childhood illnesses												
<i>Treatment of fever:</i>												
Medical treatment of fever	6.8	5.1	6.9	11.5	15.9	8.5	5.4	6.0	9.5	10.2	22.7	10.1
Treatment in a public facility	5.0	4.2	5.5	10.5	15.9	7.4	4.4	5.6	8.1	9.7	19.9	9.0
Treatment in a private facility	1.8	0.9	1.4	1.0	0.0	1.1	1.0	0.4	1.4	0.5	2.8	1.1
<i>Treatment of acute respiratory infection (ARI):</i>												
Medical treatment of ARI	22.1	13.6	13.6	19.4	33.5	19.7	6.8	14.4	20.0	28.4	50.4	23.6
Treatment in a public facility	17.7	9.1	8.6	16.5	33.5	16.1	6.8	13.2	16.1	28.4	47.4	22.0
Treatment in a private facility	4.4	4.5	5.0	2.9	0.0	3.6	0.0	1.2	3.9	0.0	3.0	1.6
<i>Treatment of diarrhea:</i>												
Use of oral rehydration therapy	37.1	37.8	42.1	48.2	63.5	43.4	39.6	49.3	35.3	49.0	69.5	46.6
Medical treatment of diarrhea	4.5	12.1	13.2	15.5	18.5	12.0	10.9	18.2	10.8	12.3	24.5	14.6
Treatment in a public facility	3.2	8.4	10.9	14.8	16.8	10.0	5.2	15.4	9.5	11.6	22.1	12.0
Treatment in a private facility	1.3	3.7	2.3	0.7	1.7	2.0	5.7	2.8	1.3	0.7	2.4	2.6
C. Contraceptive services												
<i>Contraceptive prevalence:</i>												
Women	0.5	1.4	2.0	4.2	15.3	4.5						
Men							1.0	3.9	6.5	14.3	18.7	8.4
<i>Source of contraception - public sector:</i>												
Women	*	*	(32.6)	53.9	61.8	57.2						
Men							na	na	na	na	na	na
<i>Source of contraception - private sector:</i>												
Women	*	*	(67.4)	39.0	35.6	39.3						
Men							na	na	na	na	na	na
D. Treatment of adult illnesses												
<i>Treatment of genital discharge, ulcer, sore:</i>												
Women	*	*	*	*	(65.8)	46.4						
Men							*	*	*	*	*	*
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na

Mali

1995 / 96 - FEMALE / MALE POPULATIONS

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Hygienic practices												
<i>Disposal of children's stools:</i>												
Sanitary disposal	na	na	na	na	na	na						
<i>Handwashing:</i>												
Wash hands prior to preparing food	na	na	na	na	na	na						
Handwashing facilities in household	na	na	na	na	na	na						
B. Bednet ownership and use												
<i>Bednet use:</i>												
By children	na	na	na	na	na	na						
C. Breastfeeding												
Exclusive breastfeeding	9.5	15.5	12.1	10.7	9.2	11.3	12.5	11.8	13.5	17.2	11.8	13.3
Timely complementary feeding	22.8	28.2	29.9	28.5	32.6	28.1	24.8	27.7	43.3	36.7	59.1	38.0
Bottle-feeding	1.0	0.5	2.0	5.3	10.0	3.4	2.2	1.6	1.5	2.7	7.7	3.0
D. Micronutrient consumption												
<i>Vitamin A:</i>												
Children	1.2	1.0	1.2	1.7	1.4	1.3	2.3	0.0	0.3	0.5	0.6	0.7
E. Tobacco and alcohol use												
<i>Tobacco:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
<i>Alcohol:</i>												
Women	na	na	na	na	na	na						
Men							na	na	na	na	na	na
F. Sexual practices												
<i>Non-regular sexual partnerships:</i>												
Women	na	na	na	na	na	na						
Men							8.1	10.7	16.1	20.7	39.3	19.9
<i>Condom usage with non-regular partner:</i>												
Women	na	na	na	na	na	na						
Men							6.2	13.7	18.3	32.2	52.2	35.5

Mali

1995 / 96 - FEMALE / MALE POPULATIONS

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles - Female						Wealth Quintiles - Male					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Education												
<i>School completion:</i>												
Women	1.4	2.4	3.3	8.7	36.1	10.9	4.6	6.9	9.7	19.3	52.6	20.6
Men												
<i>School participation:</i>												
Girls	5.0	10.6	14.8	26.9	55.2	22.0	11.8	14.9	21.7	33.2	62.9	28.0
Boys												
B. Exposure to mass media												
<i>Newspaper readership:</i>												
Women	0.5	0.9	1.3	4.9	21.7	6.5	3.1	4.5	6.6	14.0	44.1	15.8
Men												
<i>Radio listenership:</i>												
Women	29.1	50.6	59.0	70.2	85.3	59.8	59.0	76.8	83.1	90.3	93.9	81.3
Men												
<i>Television viewership:</i>												
Women	6.2	12.9	17.3	30.9	73.2	29.8	9.1	15.7	29.5	44.5	82.7	38.3
Men												
C. Knowledge and attitudes about HIV/AIDS												
<i>Knowledge about sexual transmission of HIV/AIDS:</i>												
Women	63.7	77.9	79.2	85.0	93.7	80.4	90.7	94.4	95.2	94.8	99.0	95.0
Men												
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>												
Women	29.6	26.7	32.6	47.2	73.5	40.8	46.6	43.5	50.0	60.1	77.3	55.7
Men												
<i>Attitudes toward HIV/AIDS:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men												
D. Orphanhood												
Paternal orphan prevalence	3.7	3.5	2.9	3.5	4.5	3.6	3.9	3.8	4.0	4.9	4.3	4.2
Maternal orphan prevalence	2.4	1.9	2.1	1.9	2.4	2.1	2.6	1.5	2.7	2.1	1.8	2.1
Double orphan prevalence	0.2	0.3	0.4	0.3	0.6	0.4	0.7	0.2	0.3	0.4	0.9	0.5

Mali
1995 / 96 - RURAL / URBAN POPULATIONS

Part I: HNP STATUS

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood illness and mortality												
Infant mortality rate	152.7	149.1	140.7	140.1	121.4	145.0	(127.0)	121.6	126.0	105.6	85.1	98.7
Under-five mortality rate	300.0	284.0	253.0	255.1	223.2	272.6	(257.4)	(279.4)	(242.6)	211.3	153.4	190.3
Prevalence of fever	40.8	40.8	40.6	40.6	38.3	40.6	66.3	49.3	36.7	35.4	28.4	33.6
Prevalence of diarrhea	28.6	26.8	28.5	27.5	21.9	27.6	38.1	27.5	22.8	22.4	14.9	19.2
Prevalence of acute respiratory infection	15.3	16.9	15.3	15.9	13.9	15.8	17.0	13.3	12.4	14.3	13.9	13.9
B. Fertility												
Total fertility rate	7.0	7.4	7.6	7.4	*	7.3	*	*	*	(6.6)	4.7	5.4
Adolescent fertility rate	197.0	224.0	216.0	241.0	*	215.6	*	*	*	(180.0)	116.0	140.2
C. Nutritional status												
<i>Children:</i>												
Moderate stunting	14.1	17.5	17.0	18.0	14.8	16.5	(16.2)	17.0	13.7	15.0	11.4	13.1
Severe stunting	18.1	18.1	15.5	15.6	9.9	16.6	(14.3)	12.5	12.0	10.3	6.5	8.7
Moderate underweight	26.4	27.6	23.2	22.0	22.9	24.9	(12.8)	25.9	21.2	19.9	18.8	19.6
Severe underweight	20.9	18.5	16.8	17.9	7.5	18.1	(31.7)	15.8	14.3	14.7	8.8	12.1
Mild anemia	na	na	na	na	na	na	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na	na	na	na	na	na	na
<i>Women:</i>												
Malnutrition	16.3	17.5	15.9	17.6	17.2	16.8	(8.2)	27.4	17.0	19.0	11.0	14.6
Mild anemia	na	na	na	na	na	na	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na	na	na	na	na	na	na
D. Female circumcision												
<i>Prevalence of circumcision:</i>												
Girls	68.1	74.5	77.0	79.7	81.5	74.7	57.0	69.8	72.8	77.3	78.5	76.4
Women	88.9	97.5	98.3	98.9	95.9	95.6	73.1	77.3	85.5	90.3	92.2	89.8
<i>Prevalence of occlusion:</i>												
Girls	na	na	na	na	na	na	na	na	na	na	na	na
Women	0.2	0.9	0.1	0.2	0.0	0.4	0.0	1.2	2.0	0.7	0.6	0.8
E. Sexually transmitted disease												
<i>Prevalence of genital discharge:</i>												
Women	0.1	0.4	0.3	0.2	0.6	0.3	0.1	0.8	0.0	0.9	0.2	0.3
Men	2.0	1.6	3.4	1.3	2.6	2.1	(3.8)	0.0	0.0	3.9	2.3	2.5
<i>Prevalence of genital ulcer:</i>												
Women	0.2	0.2	0.2	0.0	0.0	0.2	0.0	0.0	0.0	0.4	0.3	0.3
Men	2.0	3.3	0.7	3.6	1.3	2.3	(0.4)	(0.8)	0.0	2.6	0.7	1.1

Mali

1995 / 96 - RURAL / URBAN POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization												
BCG coverage	(66.8)	(87.1)	(89.4)	90.5	95.3	92.1	56.0	69.4	74.8	79.8	92.0	70.6
Measles coverage	(35.2)	(43.2)	(59.2)	62.1	80.1	69.8	34.1	41.2	47.7	47.1	78.2	43.9
DPT coverage	(17.3)	(61.5)	(51.5)	62.4	69.6	64.1	17.4	28.8	29.0	31.8	57.4	28.0
Full basic coverage	(16.4)	(42.7)	(45.6)	49.6	56.9	51.9	15.8	24.9	24.8	26.8	50.0	24.2
No basic coverage	(33.2)	(12.9)	(10.6)	8.2	4.2	7.2	43.0	30.6	24.1	17.9	8.0	28.4
Hepatitis B coverage	na	na	na	na	na	na	na	na	na	na	na	na
Yellow fever coverage	(3.2)	(1.4)	(10.5)	21.7	31.5	24.5	0.4	1.3	2.3	0.8	7.9	1.5
B. Treatment of childhood diseases												
<i>Treatment of fever:</i>												
Medical treatment of fever	3.6	4.9	13.2	10.4	24.2	16.1	6.3	5.6	7.6	11.0	5.8	7.3
Treatment in a public facility	3.6	2.2	13.2	10.3	22.1	14.9	4.8	5.1	6.0	10.0	5.8	6.2
Treatment in a private facility	0.0	2.7	0.0	0.1	2.1	1.2	1.5	0.5	1.6	1.0	0.0	1.1
<i>Treatment of acute respiratory infection (ARI):</i>												
Medical treatment of ARI	*	*	*	27.3	49.9	37.5	15.9	14.2	16.3	22.3	*	16.6
Treatment in a public facility	*	*	*	24.7	47.9	35.1	13.3	11.8	11.3	21.4	*	13.9
Treatment in a private facility	*	*	*	2.6	2.0	2.4	2.6	2.4	5.0	0.9	*	2.7
<i>Treatment of diarrhea:</i>												
Use of oral rehydration therapy	39.5	(34.8)	44.6	54.1	75.0	59.2	38.3	44.3	37.9	46.2	(44.4)	41.4
Medical treatment of diarrhea	14.2	(25.9)	13.1	15.8	26.1	20.6	7.3	14.3	11.9	13.3	(10.2)	11.5
Treatment in a public facility	14.2	(21.1)	13.1	14.5	23.3	18.6	3.6	11.2	9.9	12.8	(10.2)	9.1
Treatment in a private facility	0.0	(4.8)	0.0	1.3	2.8	2.0	3.7	3.1	2.0	0.5	0.0	2.4
C. Antenatal and delivery care												
<i>Antenatal care visits:</i>												
To a medically trained person	35.3	49.2	68.3	81.4	90.0	81.0	19.9	34.0	40.8	46.3	68.1	35.8
To a doctor	0.4	0.9	1.2	2.1	3.5	2.6	0.1	0.6	0.7	1.0	0.6	0.6
To a nurse or trained midwife	34.9	48.3	67.1	79.2	86.5	78.4	19.7	33.4	40.1	45.3	67.5	35.2
Multiple visits to a medically trained person	25.7	22.9	50.4	65.5	74.8	64.9	10.1	21.2	29.1	33.0	48.3	23.6
<i>Antenatal care content:</i>												
Tetanus toxoid	42.5	53.9	67.6	79.3	85.9	78.8	30.7	40.1	44.7	50.1	66.4	41.8
Prophylactic antimalarial treatment	na	na	na	na	na	na	na	na	na	na	na	na
Iron supplementation	na	na	na	na	na	na	na	na	na	na	na	na
<i>Delivery attendance:</i>												
By a medically trained person	32.4	46.9	60.6	79.8	88.6	79.0	10.1	20.9	30.5	37.5	48.5	24.9
By a doctor	0.0	0.0	1.4	1.9	2.2	1.9	0.1	0.3	0.1	0.8	0.9	0.3
By a nurse or trained midwife	32.4	46.9	59.1	77.9	86.3	77.1	10.0	20.6	30.4	36.6	47.6	24.5
In a public facility	32.1	46.9	51.0	70.7	75.2	68.4	7.3	13.6	19.8	24.7	36.9	16.6
In a private facility	0.0	0.0	0.0	0.3	2.8	1.6	0.1	0.0	0.3	0.2	0.0	0.1
At home	67.9	52.9	39.6	22.6	11.2	21.6	90.7	80.6	70.9	65.3	55.2	76.8
D. Contraceptive services												
<i>Contraceptive prevalence:</i>												
Women	1.5	0.2	5.8	6.1	17.0	11.6	0.4	1.5	1.5	3.3	8.9	1.9
Men	(4.9)	(5.3)	0.0	15.9	19.2	15.1	0.8	3.8	7.6	13.2	16.2	5.8

Mali

1995 / 96 - RURAL / URBAN POPULATIONS

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
D. Contraceptive services (cont.)												
<i>Source of contraception - public sector:</i>												
Women	*	*	*	(74.7)	62.5	62.9	*	*	*	(35.1)	(57.3)	44.9
Men	na	na	na	na	na	na	na	na	na	na	na	na
<i>Source of contraception - private sector:</i>												
Women	*	*	*	(21.9)	35.1	34.7	*	*	*	(54.4)	(39.4)	49.4
Men	na	na	na	na	na	na	na	na	na	na	na	na
E. Treatment of adult illnesses												
<i>Treatment of genital discharge, ulcer, sore:</i>												
Women	*	*	*	*	*	(66.3)	*	*	*	*	*	(29.5)
Men	*	*	*	*	*	(42.9)	*	*	*	*	*	26.4
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
<i>Voluntary counseling and testing for HIV/AIDS:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na

Mali

1995 / 96 - RURAL / URBAN POPULATIONS

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Hygienic practices												
<i>Disposal of children's stools:</i>												
Sanitary disposal	na	na	na	na	na	na	na	na	na	na	na	na
<i>Handwashing:</i>												
Wash hands prior to preparing food	na	na	na	na	na	na	na	na	na	na	na	na
Handwashing facilities in household	na	na	na	na	na	na	na	na	na	na	na	na
B. Bednet ownership and use												
<i>Bednet ownership:</i>												
Bednet ownership	na	na	na	na	na	na	na	na	na	na	na	na
Treated bednet ownership	na	na	na	na	na	na	na	na	na	na	na	na
<i>Bednet use:</i>												
By children	na	na	na	na	na	na	na	na	na	na	na	na
By pregnant women	na	na	na	na	na	na	na	na	na	na	na	na
C. Breastfeeding												
Exclusive breastfeeding	*	*	*	19.7	11.9	16.5	10.8	11.9	10.9	10.3	*	10.7
Timely complementary feeding	*	*	*	(46.2)	49.4	45.1	23.5	27.6	36.8	25.6	*	29.0
Bottle-feeding	0.0	0.6	4.4	7.5	10.2	7.8	1.6	1.1	1.4	2.0	4.1	1.6
D. Micronutrient consumption												
<i>Iodized salt:</i>												
Availability of iodized salt in household	na	na	na	na	na	na	na	na	na	na	na	na
<i>Vitamin A:</i>												
Children	0.0	2.0	2.6	0.8	1.3	1.3	1.8	0.4	0.5	1.2	0.0	0.9
Women	na	na	na	na	na	na	na	na	na	na	na	na
E. Tobacco and alcohol use												
<i>Tobacco:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
<i>Alcohol:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
F. Sexual practices												
<i>Non-regular sexual partnerships:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	14.7	4.3	22.1	24.7	42.7	34.2	7.5	11.1	15.1	18.3	19.9	12.9
<i>Condom usage with non-regular partner:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	23.4	8.5	44.8	48.9	53.6	51.6	3.4	13.8	11.8	18.3	35.7	14.6
G. Domestic violence												
Ever experienced violence	na	na	na	na	na	na	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na	na	na	na	na	na	na

Mali

1995 / 96 - RURAL / URBAN POPULATIONS

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles - Rural						Wealth Quintiles - Urban					
	Low	2nd	3rd	4th	High	Avg.	Low	2nd	3rd	4th	High	Avg.
A. Education												
<i>School completion:</i>												
Women	3.3	7.2	4.4	15.2	40.4	28.4	1.3	2.0	3.1	5.2	15.0	3.3
Men	8.6	10.6	15.1	28.7	54.8	42.5	4.4	6.6	8.8	13.5	38.9	9.5
<i>School participation:</i>												
Girls	1.2	16.7	28.4	38.6	60.1	47.1	5.2	9.9	12.9	21.6	33.6	12.6
Boys	11.6	20.1	32.4	48.1	67.5	52.9	11.8	14.3	20.1	26.3	46.8	19.2
B. Exposure to mass media												
<i>Newspaper readership:</i>												
Women	1.7	3.8	2.9	8.6	24.3	17.2	0.4	0.6	1.1	2.9	8.1	1.5
Men	7.4	9.6	19.2	20.0	47.8	35.9	2.7	4.2	4.5	10.4	22.8	6.0
<i>Radio listenership:</i>												
Women	28.0	56.8	65.4	78.1	87.7	80.0	29.1	50.0	58.0	65.8	72.5	50.4
Men	55.9	78.8	82.7	95.5	95.1	91.8	59.3	76.7	83.2	87.2	87.0	76.1
<i>Television viewership:</i>												
Women	4.8	15.1	26.7	41.0	76.6	58.8	6.3	12.7	15.7	25.2	55.2	16.3
Men	6.2	15.2	51.2	59.7	87.5	71.8	9.4	15.8	25.9	35.1	54.5	21.9
C. Knowledge and attitudes about HIV/AIDS												
<i>Knowledge about sexual transmission of HIV/AIDS:</i>												
Women	59.2	74.8	73.4	89.8	95.0	89.7	64.0	78.2	80.1	82.4	87.0	76.1
Men	75.8	89.9	92.9	98.5	99.6	97.4	92.0	94.7	95.5	92.6	95.7	93.8
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>												
Women	20.9	27.8	40.3	58.8	76.8	65.2	17.6	26.6	31.4	40.7	56.5	29.5
Men	37.1	41.5	45.4	68.9	78.8	70.6	42.7	43.7	50.8	54.8	68.4	48.4
<i>Attitudes toward HIV/AIDS:</i>												
Women	na	na	na	na	na	na	na	na	na	na	na	na
Men	na	na	na	na	na	na	na	na	na	na	na	na
D. Status of women												
<i>Household decisionmaking:</i>												
Can seek own health care	na	na	na	na	na	na	na	na	na	na	na	na
Can seek children's health care	na	na	na	na	na	na	na	na	na	na	na	na
Can make daily household purchases	na	na	na	na	na	na	na	na	na	na	na	na
Can make large household purchases	na	na	na	na	na	na	na	na	na	na	na	na
Can make meal-related decisions	na	na	na	na	na	na	na	na	na	na	na	na
<i>Freedom of movement:</i>												
Can travel to visit family, relatives	na	na	na	na	na	na	na	na	na	na	na	na
<i>Other decisionmaking, attitudes:</i>												
Can decide how to spend own money	na	na	na	na	na	na	na	na	na	na	na	na
Can decide whether to have sex	na	na	na	na	na	na	na	na	na	na	na	na
Justifies domestic violence	na	na	na	na	na	na	na	na	na	na	na	na
E. Orphanhood												
Paternal orphan prevalence	7.8	6.0	4.9	6.4	5.3	5.6	3.6	3.4	3.2	3.0	1.2	3.2
Maternal orphan prevalence	3.9	1.9	3.2	1.7	2.2	2.2	2.4	1.7	2.3	2.2	1.7	2.1
Double orphan prevalence	1.6	0.4	0.4	0.3	0.8	0.7	0.4	0.3	0.3	0.4	0.3	0.3

PART III. TECHNICAL NOTES

- A. INDICATOR DEFINITIONS**
- B. DATA AND METHODS**
- C. DISCUSSION**

INDICATOR DEFINITIONS

Part I: HNP STATUS

A. CHILDHOOD MORTALITY AND ILLNESS ²

Infant mortality rate: number of deaths to children under 12 months of age per 1,000 live births, based on experience during the ten years preceding the survey.

Under-five mortality rate: number of deaths to children under five years of age per 1,000 live births, based on experience during the ten years preceding the survey.

Prevalence of fever: percent of children who had fever, whether or not accompanied by cough or rapid breathing, in the two weeks before the survey.

Prevalence of diarrhea: percent of children who had diarrhea in the two weeks before the survey.

Prevalence of acute respiratory infection: percent of children who had a cough accompanied by rapid or difficult breathing in the two weeks before the survey.

B. FERTILITY

Total fertility rate (TFR): average number of births a woman could expect to have during her lifetime if she followed the levels of fertility currently observed at every age. The TFR is calculated as the sum of average annual age-specific fertility rates for all reproductive age groups (usually 15-49 years) in the three years before the survey.

Adolescent fertility rate: age-specific fertility rate for women 15-19 years of age. This is the average number of births among women aged 15-19 years per 1,000 women in that age group, based on births in the three years before the survey and expressed as annual averages.

² Figures for the prevalence of fever, diarrhea, and acute respiratory infection refer to percent of children under three, four, or five years of age, depending upon the country. (The specific ages covered for in particular country may be determined by consulting the full report on that country's DHS, which may be found at: www.measuredhs.com/countries.)

C. NUTRITIONAL STATUS

Children^{3, 4, 5}

Moderate stunting (height-for-age): percent of children with a height-for-age Z-score of between –2 and –3 standard deviations of the median reference standard for their age (as defined in fn. 4).

Severe stunting (height-for-age): percent of children with a height-for-age Z-score of below –3 standard deviations of the median reference standard for their age (as defined in fn. 4).

Moderate underweight (weight-for-age): percent of children with a weight-for-age Z-score of between –2 and –3 standard deviations of the median reference standard for their age (as defined in fn. 4).

Severe underweight (weight-for-age): percent of children with a weight-for-age Z-score of below –3 standard deviations of the median reference standard for their age (as defined in fn. 4).

Mild anemia: percent of children with a hemoglobin level of between 10.0g/dl and 10.9 g/dl, the World Health Organization criterion for mild anemia.

Moderate anemia: percent of children with a hemoglobin level of between 7.0g/dl and 9.9g/dl, the World Health Organization criterion for moderate anemia.

Severe anemia: percent of children with a hemoglobin level of below 7.0g/dl, the World Health Organization criterion for severe anemia.

Women^{6, 7}

Malnutrition: percent of women aged 15-49 years with a Body Mass Index (BMI) of less than 18.5, where BMI – commonly used to indicate adult nutritional status – is defined as weight in kilograms divided by the square of height in meters.

Mild anemia: percent of women aged 15-49 years with a hemoglobin level of between 10.0g/dl and 10.9g/dl for pregnant women and between 10.0g/dl and 11.9g/dl for non-pregnant women, the World Health Organization criterion for mild anemia.

Moderate anemia: percent of women aged 15-49 years with a hemoglobin level of between 7.0g/dl and 9.9g/dl, the World Health Organization criterion for moderate anemia.

Severe anemia: percent of women aged 15-49 years with a hemoglobin level of less than 7.0g/dl, the World Health Organization criterion for severe anemia.

³ All figures related to children's nutrition status refer to children under three, four, or five years of age, depending upon the country. (The specific ages covered for in particular country may be determined by consulting the full report on the DHS of the country concerned, which is available at: www.measuredhs.com/countries.)

⁴ The reference standards used for stunting and underweight are those established in the 1970s by the World Health Organization, the U.S. Centers for Disease Control, and the U.S. National Center for Health Statistics. Updated stunting and underweight figures based on the recently-released, 2006 World Health Organization reference standards are currently under preparation. When complete, they will be available at: www.worldbank.org/povertyandhealth/countrydata.

⁵ The anemia figures for children living at an altitude above 1,000 meters have been adjusted to reflect the higher level of hemoglobin required.

⁶ In some countries, the BMI is presented for all women; in others, the figure is available only for mothers of children under five years of age. The reference population for any given country can be determined by consulting the full report on the DHS for the country concerned. An electronic version of this report is located at: www.measuredhs.com/countries.

⁷ Anemia cut-off points for respondents who live at an altitude above 1,000 meters and/or who smoke have been adjusted to account for their higher hemoglobin requirements.

D. FEMALE CIRCUMCISION

Prevalence of Circumcision

Girls: percent of women aged 15-49 years with one or more daughters, at least one of whom had been circumcised.

Women: percent of women aged 15-49 years who had been circumcised.

Prevalence of Occlusion

Girls: percent of women aged 15-49 years with one or more daughters, at least one of whom had been circumcised with the vaginal area sewn closed.

Women: percent of women aged 15-49 years whose vaginal area had been sewn closed.

E. SEXUALLY TRANSMITTED DISEASES

Prevalence of Genital Discharge

Women: percent of women aged 15-49 years who had had abnormal genital discharge in the twelve months before the survey.

Men: percent of men aged 15-54 years who had had abnormal genital discharge in the twelve months before the survey.

Prevalence of Genital Ulcer or Sore

Women: percent of women aged 15-49 years who had had a genital ulcer or sore in the twelve months before the survey.

Men: percent of men aged 15-54 years who had had a genital ulcer or sore in the twelve months before the survey.

Part II: INTERMEDIATE DETERMINANTS OF HNP STATUS – HEALTH SERVICE USE

A. CHILDHOOD IMMUNIZATION ⁸

BCG coverage: percent of children who had received a dose of BCG vaccine by the time of the survey.

Measles coverage: percent of children who had received a dose of measles vaccine by the time of the survey.

DPT coverage: percent of children who had received three doses of DPT vaccine by the time of the survey.

Full basic coverage: percent of children who had received a dose of BCG vaccine, measles vaccine, and three doses of DPT and polio vaccines by the time of the survey, excluding polio vaccine given at birth.

No basic coverage: percent of children who had received no vaccination against the six early-childhood diseases (TB, measles, polio, diphtheria, pertusis, and tetanus) by the time of the survey.

Hepatitis B coverage: percent of children who had received three doses of hepatitis B vaccine by the time of the survey.

Yellow fever coverage: percent of children who had received a dose of yellow fever vaccine by the time of the survey.

B. TREATMENT OF CHILDHOOD ILLNESSES ⁹

Treatment of Fever

Medical treatment of fever: percent of children with fever, with or without cough or rapid breathing, in the two weeks before the survey who had sought medical advice for fever from any health facility or health provider, whether public or private.

Treatment in a public facility: percent of children with fever, with or without cough or rapid breathing, in the two weeks before the survey who had sought medical advice for fever from a public-sector health facility or provider (as defined in fn. 9).

Treatment in a private facility: percent of children with fever, with or without cough or rapid breathing, in the two weeks before the survey who had sought medical advice for fever from a private-sector health facility or provider (as defined in fn. 9).

⁸ Childhood immunization figures refer to rates among children 12-23 months of age in all countries except those in Latin America and the Caribbean. There, the figures refer to rates among children 18-29 months of age. All figures are based on information recorded on the child's vaccination card; or, in cases where a card was not seen by the interviewer, on the mother's report.

⁹ Figures for illness treatment in a public facility refer to treatment in government hospitals, health centers, health posts, or dispensaries; or in facilities operated by government-affiliated social securing programs. Figures for treatment in private facilities cover treatment in private hospitals or clinics, in private doctors' offices, or in facilities operated by other private medical providers (such as non-governmental organizations) as defined in the country concerned; but exclude treatment obtained in private pharmacies or shops.

Treatment of Acute Respiratory Infection (ARI)

Medical treatment of ARI: percent of children with a cough and rapid breathing in the two weeks before the survey who had been taken for treatment at any medical facility or provider, whether public or private.

Treatment in a public facility: percent of children with a cough and rapid breathing in the two weeks before the survey who had been taken for treatment at a public-sector health facility or provider (as defined in fn. 9).

Treatment in a private facility: percent of children with a cough and rapid breathing in the two weeks before the survey who had been taken for treatment at a private-sector health facility or provider (as defined in fn. 9).

Treatment of Diarrhea

Use of oral rehydration therapy: percent of children with diarrhea in the two weeks before the survey who had received oral rehydration therapy (ORT) (defined as including consumption of oral rehydration salts, other recommended home fluids, or other increased liquids).

Medical treatment of diarrhea: percent of children with diarrhea in the two weeks before the survey who had been taken for treatment at any medical facility or provider, whether public or private.

Treatment in a public facility: percent of children with diarrhea in the two weeks before the survey who had been taken for treatment at a public-sector health facility or provider (as defined in fn. 9).

Treatment in a private facility: percent of children with diarrhea in the two weeks before the survey who had been taken for treatment at a private-sector health facility or provider (as defined in fn. 9).

C. ANTENATAL AND DELIVERY CARE ¹⁰

Antenatal Care (ANC) Visits

To a medically-trained person: percent of women with one or more births in the five years before the survey who had received at least one antenatal care consultation from a medically-trained person (as defined in fn. 10) before her most recent birth.

To a doctor: percent of women with one or more births in the five years before the survey who had received at least one antenatal care consultation from a doctor before her most recent birth.

To a nurse or trained midwife: percent of women with one or more births in the five years before the survey who had received at least one antenatal care consultation from a nurse or trained midwife (as defined in fn. 10) before her most recent birth.

Multiple visits to a medically-trained person: percent of women with one or more births in the five years before the survey who had received at least three antenatal care consultations from any medically-trained provider (as defined in fn. 10) before her most recent birth.

¹⁰ When speaking of antenatal and delivery care, medically-trained persons are defined as doctors, nurses, and trained midwives. The definition excludes traditional midwives or other traditional birth attendants, whether trained or untrained.

Antenatal Care (ANC) Content

Tetanus toxoid: percent of women with one or more births in the five years before the survey who had received at least one tetanus toxoid injection during her most recent pregnancy.

Prophylactic antimalarial treatment: percent of women with one or more births in the five years before the survey who had received prophylactic treatment with any anti-malarial drug during her most recent pregnancy.

Iron supplementation: percent of women with one or more births in the five years before the survey who had taken iron tablets during her most recent pregnancy.

Delivery Attendance

By a medically-trained person: percent of births in the five years before the survey attended by a medically-trained person (as defined in fn. 10).

By a doctor: percent of births in the five years before the survey attended by a doctor.

By a nurse or trained midwife: percent of births in the five years before the survey attended to by a nurse or a trained midwife (as defined in fn. 10).

In a public facility: percent of all deliveries in the five years before the survey occurring in a public-sector health facility (as defined in fn. 9).

In a private facility: percent of all deliveries in the five years before the survey occurring in a private-sector health facility (as defined in fn. 9).

At home: percent of all deliveries in the five years before the survey occurring at home (defined as the woman's own or any other home).

D. CONTRACEPTIVE SERVICES

Contraceptive Prevalence¹¹

Women: percent of married or in-union women aged 15-49 years who used any modern means of contraception (as defined in fn. 11).

Men: percent of married or in-union men aged 15-54 years who used any modern means of contraception (as defined in fn. 11).

Source of Contraception - Public Sector

Women: percent of married women who obtained their current method of contraception from a public-sector health facility or provider (as defined in fn. 9).

Men: percent of married men who obtained their current method of contraception from a public-sector health facility or provider (as defined in fn. 9).

¹¹ Figures refer to use of modern means of contraception, defined as male/female sterilization, oral contraceptive pill, contraceptive injection, intrauterine device, male/female condom, diaphragm, cervical cap, contraceptive jelly or foam, implant, or some country-specific modern method.

Source of Contraception - Private Sector

Women: percent of married women who obtained their current method of contraception from a private-sector health facility or provider (as defined in fn. 9, except that private pharmacies and shops are included rather than excluded).

Men: percent of married women who obtained their current method of contraception from a private-sector health facility or provider (as defined in fn. 9, except that private pharmacies and shops are included rather than excluded).

E. TREATMENT OF ADULT ILLNESSES

Treatment of Genital Discharge, Ulcer, or Sore

Women: percent of women with genital discharge, ulcer, or sore who sought any medical treatment for resulting symptoms.

Men: percent of men with genital discharge, ulcer, or sore who sought any medical treatment for resulting symptoms.

Treatment of Genital Discharge, Ulcer, or Sore in a Public Facility

Women: percent of women with genital discharge, ulcer, or sore who sought treatment from a public-sector health facility or provider (as defined in fn. 9).

Men: percent of men with genital discharge, ulcer, or sore who sought treatment from a public-sector health facility or provider (as defined in fn. 9).

Voluntary Counseling and Testing for HIV/AIDS

Women: percent of women aged 15-49 years who had been tested for HIV at any time before the survey.

Men: percent of men aged 15-54 years who had been tested for HIV at any time before the survey.

Part III: INTERMEDIATE DETERMINANTS OF HNP STATUS – INDIVIDUAL AND HOUSEHOLD BEHAVIOR

A. HYGIENIC PRACTICES

Disposal of Children’s Stools

Sanitary disposal: percent of mothers with at least one child under five years of age who disposed of the stools of their youngest child in a sanitary manner (defined as dropping stool into a latrine, burying it, or using disposable diapers).

Handwashing

Wash hands prior to preparing food: percent of women aged 15-49 years preparing meals who washed hands before handling food.

Handwashing facilities in household: percent of households that had hand-washing materials or facilities, as determined by direct observation of interviewers.

B. BEDNET OWNERSHIP AND USE

Bednet Ownership

Bednet ownership: percent of households owning one or more bednets.

Treated bednet ownership: percent of households owning one or more bednets that had recently been treated with insecticides.

Bednet Use

By children: percent of households with at least one child under five years of age, some or all of whom had slept under a bednet the night before the survey.

By pregnant women: percent of currently pregnant women who had slept under a bednet the night before the survey.

C. BREASTFEEDING

Exclusive breastfeeding: percent of children 0-3 months of age who had received only breast milk in the 24 hours before the survey.

Timely complementary feeding: percent of children 6-9 months of age who had received breast milk and solid or semi-solid foods in the twenty-four hours before the survey.

Bottle-feeding: percent of children under 12 months of age who had received any food or drink from a bottle with a nipple in the twenty-four hours before the survey.

D. MICRONUTRIENT CONSUMPTION

Iodized Salt

Availability of iodized salt in household: percent of households with cooking salt testing positive for iodine/iodate at the recommended level of 15 or 25 parts per million or more (depending on the country).¹²

Vitamin A

Children: percent of children¹³ who had received at least one dose of vitamin A in the six months before the survey, as reported by the mothers.

Women: percent of women who had received a dose of vitamin A within two months of the last birth, in the five years before the survey.

E. TOBACCO AND ALCOHOL USE

Tobacco¹⁴

Women: percent of women aged 15-49 years who currently were smoking or chewing tobacco products.

Men: percent of men aged 15-54 years who currently were smoking or chewing tobacco products.

Alcohol

Women: percent of women aged 15-49 years who had gotten intoxicated due to excessive consumption of alcohol in the three months before the survey.

Men: percent of men aged 15-54 years who had gotten intoxicated due to excessive consumption of alcohol in the three months before the survey.

F. SEXUAL PRACTICES

Non-Regular Sexual Partnerships

Women: percent of women aged 15-49 years who had had sex with a non-regular partner at least once in the twelve months before the survey.

Men: percent of men aged 15-54 years who had had sex with a non-regular partner at least once in the twelve months before the survey.

¹² Which of these two levels is recommended in any particular country may be determined by consulting the full report on that country's DHS, which may be found at: www.measuredhs.com/countries.

¹³ Figures refer to children over six months of age and under three, four, or five years of age, depending upon the country. (The specific ages covered for in a particular country may be determined by consulting the full report on that country's DHS, which is available at: www.measuredhs.com/countries.)

¹⁴ Tobacco products include cigarettes, pipes, cigars, leaves, etc.

Condom Usage with Non-Regular Partner

Women: percent of women aged 15-49 years with non-regular sexual partner who had used a condom in the last sexual intercourse with such a partner in the twelve months before the survey.

Men: percent of men aged 15-54 years with non-regular sexual partner who had used a condom in the last sexual intercourse with such a partner in the twelve months before the survey.

G. DOMESTIC VIOLENCE

Ever experienced violence: percent of women aged 15-49 years who had ever been hit or beaten by current or former husband/partner.

Experienced violence in past year: percent of women aged 15-49 years who had been hit or beaten by current or former husband/partner in the twelve months before the survey.

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

A. EDUCATION

School Completion

Women: percent of women aged 15-49 years who had completed the fifth grade.

Men: percent of men aged 15-54 years who had completed the fifth grade.

School Participation

Girls: percent of girls aged 6-10 years who were attending school at the time of the survey.

Boys: percent of boys aged 6-10 years who were attending school at the time of the survey.

B. EXPOSURE TO MASS MEDIA

Newspaper Readership

Women: percent of women aged 15-49 years who read a newspaper at least once a week.

Men: percent of men aged 15-54 years who read a newspaper at least once a week.

Radio Listenership

Women: percent of women aged 15-49 years who listened to radio at least once a week.

Men: percent of men aged 15-54 years who listened to radio at least once a week.

Television Viewership

Women: percent of women aged 15-49 years who watched television at least once a week.

Men: percent of men aged 15-54 years who watched television at least once a week.

C. KNOWLEDGE AND ATTITUDES ABOUT HIV/AIDS

Knowledge about Sexual Transmission of HIV/AIDS¹⁵

Women: percent of women aged 15-49 years who knew of HIV/AIDS and of at least one of the following ways to avoid it through interruption of its sexual transmission route: abstinence; using a condom; avoiding multiple sex partners, sex with prostitutes, and unprotected homosexual sex.

Men: percent of men aged 15-54 years who knew of HIV/AIDS and of at least one of the ways to avoid HIV/AIDS referred to in the preceding definition.

¹⁵ In most countries, the survey sample included both married and unmarried individuals. Where this was the case, all respondents, regardless of marital status, were asked the question covered in this section. Where the survey covered only individuals who were or had been married, the data pertain only to individuals who had ever been married. (The marital status of people covered for in particular country is indicated in the full report on that country's DHS, which is located at: www.measuredhs.com/countries.)

Knowledge about Mother-to-Child Transmission of HIV/AIDS

Women: percent of women aged 15-49 years who knew of at least one way HIV/AIDS can be transmitted from mother to child during pregnancy, delivery, or breastfeeding.

Men: percent of men aged 15-54 years who knew of at least one way HIV/AIDS can be transmitted from mother to child during pregnancy, delivery, or breastfeeding.

Attitudes toward HIV/AIDS

Women: percent of women aged 15-49 years who believed that people with HIV/AIDS should be allowed to continue working or that HIV test results should remain confidential.

Men: percent of men aged 15-54 years who believed that people with HIV/AIDS should be allowed to continue working or that HIV test results should remain confidential.

D. STATUS OF WOMEN

Household Decisionmaking

Can seek own health care: percent of women age 15-49 years who could decide by themselves to seek their own health care.

Can seek children's health care: percent of women aged 15-49 years, whose children live with them, who could decide by themselves to seek health care for their children.

Can make daily household purchases: percent of women aged 15-49 years who could decide by themselves or jointly with others to make daily household purchases.

Can make large household purchases: percent of women aged 15-49 years who could decide by themselves or jointly with others to make large household purchases.

Can make meal-related decisions: percent of women aged 15-49 years who could decide by themselves what food to cook daily.

Freedom of Movement

Can travel to visit family, relatives: percent of women aged 15-49 years who could decide by themselves to visit family and relatives.

Other Decisionmaking, Attitudes

Can decide how to spend own money: percent of women aged 15-49 years who work for cash who could decide by themselves on how to use the money they earn.

Can decide whether to have sex: percent of women aged 15-49 years agreeing that they can refuse to have sex with their husband for at least one of the following reasons: he has a sexually-transmitted disease; he has had sexual relations with another woman; or the woman is tired, not in mood, or recently has given birth.

Justify domestic violence: percent of women aged 15-49 years believing that a husband/male partner would be justified in beating his wife/female partner for at least one of the following reasons: he suspects her of being unfaithful; she argues with him; she goes out without telling him; she neglects the children; she burns the food; or other, country-specific reasons (for example, she shows disrespect for her in-laws or her family does not give the expected dowry).

E. ORPHANHOOD

Maternal orphan prevalence: percent of children under 15 years of age whose natural mother had died before the survey.

Paternal orphan prevalence: percent of children under 15 years of age whose natural father had died before the survey.

Double orphan prevalence: percent of children under 15 years of age both of whose natural parents had died before the survey.

DATA AND METHODS

Any assessment of the figures featured in this report requires an appreciation of how they were prepared. The first need is to understand the basic features of the data and methods employed.

A. SOURCE OF FIGURES

The figures appearing in this report are all derived from data collected under the Demographic and Health Surveys (DHS) program conducted by ORC Macro, with support from the U.S. Agency for International Development and other external assistance organizations. Large DHS household surveys have been carried out, usually at periodic intervals, in approximately seventy-five countries across Africa, Asia, Latin America, the Middle East, and the former Soviet Union.¹⁶ This series of reports covers the fifty-six of those countries that had one or more DHS surveys undertaken since 1990, for which data were publicly available as of June 2006. (Annex C is a list of the countries for which reports have been prepared.)

In each country, the DHS program has gathered information on a large number of indicators about health, nutrition, and population (hnp) status and service use; about relevant behaviors of household members; and about household characteristics like those described below. It has done this through a set of questionnaires, similar in all countries, to collect data at the individual, household, and community levels.

The data presented here draw on responses to the individual and household questionnaires. In most cases, they are based on responses from women or other family members interviewed. The principal exceptions concern nutritional status, which is based on anthropometric measurement; immunization, which typically relies to the extent possible on record cards maintained at the household level; and those other items where a source other than interviewer response is specifically identified.

B. MEASUREMENT OF ECONOMIC STATUS

Wealth or Asset Approach

Economic status has been expressed in terms of wealth or assets: specifically, on the basis of information about household characteristics gathered through the DHS household questionnaire. (Such information was normally provided for at least 25-30, and often many more, characteristics like the presence, availability, or use of a fan, radio receiver, or automobile; housing materials like wood or concrete flooring, tile or tin roofing, or cement block walls; superior sources of water like piped or a protected well; and other attributes related to economic status.)

¹⁶ Further information about the DHS program is available at the program's website: www.measuredhs.com.

Index Construction

A single, consolidated index of living standards¹⁷ was constructed by using principal components analysis (PCA) to generate a weight for each household item with available information. A wealth index score was calculated for each household by weighting the response with respect to each item pertaining to that household by the coefficient of the first principal component as determined by application of principal components analysis, and summing the results. The resulting household scores were standardized in relation to a standard normal distribution with a mean of zero and a standard deviation of one.

All individuals usually present in each household were assigned the household's standardized wealth index score, and all individuals in the sample population were ranked according to that score. The sample population was then divided into quintiles of individuals, with all individuals in a single household being assigned to the same quintile.

The same standardized household wealth index scores originally derived for the total population sample, as just described, were also used in preparing the disaggregated estimates for female and male members of the sample population, and for rural and urban residents. In preparing those disaggregated estimates, the entire population sample was divided into quintiles of individuals; the females and males, and the rural and urban residents in each quintile of the entire sample were then separated from one another; and the mean for each of the ten resulting subgroups (five female, five male; or five rural, five urban) was calculated.

This procedure was carried out separately for each of the surveys covered.

C. CALCULATION AND PRESENTATION OF RATES

Use of Sampling Weights

Rates for all health, nutrition, and population indicators are calculated after applying the DHS sampling weights. (DHS surveys often over-sample certain small subgroups of interest – residents of a particular geographic area, for example – in order to get sample sizes large enough to produce statistically-significant results. The DHS sampling weights are used to compensate for such over-sampling in order to ensure that the results are representative of the population as a whole and not just of the DHS sample.)

Calculation of Total Population Averages

The average for the total population presented alongside the quintile-specific rates for each indicator is calculated without reference to quintile divisions. It thus equals the weighted mean of the quintile rates, with the weight assigned to each quintile rate being the proportion of the number of individuals at risk (as defined on p. 59) for the indicator concerned.

Sampling Errors

Information needed to assess the statistical significance of differences among the quintile-specific rates is presented in three ways:

- First, in all the basic tables presented in part I, rates are shown in parentheses or replaced by asterisks in cases where the standard error is likely to be unacceptably high because

¹⁷ Such an index is usually referred to as either an “asset index” or a “wealth index.” The two expressions are used interchangeably in this report; for ease of communication, “wealth index” appears more frequently despite the inexact correspondence between the items included in the index's construction and those appearing in more conventional, financially-based definitions of wealth.

of small sample size. The number of observations used to determine how to present the data for the different indicators covered were as follows:

Indicator	Unit of measure	Number of observations used to determine how quintile-specific rate was presented		
		<i>Without parentheses</i>	<i>With parentheses</i>	<i>Replaced by asterisk</i>
Infant and child mortality	Number of deaths	>500	250-499	<250
Total and adolescent fertility	Number of births	>250	125-249	<125
All other indicators	Number of individuals	>50	25-49	<25

- Second, the standard error for each quintile-specific rate (except for any rate replaced by an asterisk) appearing in the total population table is provided in part III.B. (Standard errors for the quintile-specific rates presented in the female-male and rural-urban tables are not available.)

- Third, the right-hand column of the total population table provides the standard error for the concentration index, one of the measures of inequality shown, as indicated below.

D. MEASUREMENT OF INEQUALITY

Accompanying each of the indicators presented in the total population table are the values for three statistical measures of inequality:

- *Low/High Quintile Ratio*: the ratio between the rate prevailing in the lowest (poorest) population quintile and that found in the highest (least poor) quintile.

- *Low-High Quintile Difference*: the value of the lowest quintile minus the value of the highest, expressed as an absolute value.

- *Concentration Index*: twice the area in a Lorenz-type diagram between the line of equality and the concentration curve for the indicator in question, the curve being the graph of the cumulative share of the indicator against the cumulative share in the asset distribution. (The value, which can range from -1 to +1, is negative when the hnp indicator is higher among the poor (e.g., fertility), positive when it is higher among the better-off (e.g., modern contraceptive use), and zero when on balance the indicator shows no systematic relationship with wealth.)¹⁸

¹⁸ Adam Wagstaff, Pierella Paci, and Eddy van Doorslaer, "On the Measurement of Inequalities of Health," *Social Science and Medicine* 33 (1991): 545-57. (See also chapter eight in the volume by O'Donnell, van Doorslaer, Wagstaff, and Lindelow described in the annotated bibliography that constitutes annex A.)

DISCUSSION

While a basic understanding of the data and methods employed is necessary to adequately appreciate the figures appearing in this report, it is not sufficient. For the application of the approach taken involves many subtleties that also need to be understood. Among the more important are:

A. DESCRIPTIVE NATURE OF THE RELATIONSHIPS

The hnp-poverty relationships shown in this report are no more than descriptive. They should not be taken to imply any direct causal relationships, for several reasons.

One reason is the possibility that it is not wealth or asset possession *per se* that determine a person's health condition. Rather, the determining factors could be other characteristics (such as education or ethnic background) that are simultaneously associated with both asset ownership and health status.

It is also possible that the health-poverty relationships shown are driven primarily by particular items included in the index (e.g., water and sanitation). Should this be the case, improvements in health conditions among the poor might be more effectively brought about by focusing on changing those particular components of the wealth index rather than by a general effort to increase economic status as measured by the index as a whole.

B. IMPLICATIONS OF A WEALTH/ASSET APPROACH

Wealth or Assets as a Measure of Economic Status

Reliance on a wealth index to measure economic status is a rather recent development in research on economic disparities, where such status traditionally has been defined in terms of consumption or income. The principal reason for the choice of the wealth index is pragmatic rather than conceptual: the DHS surveys, which are of interest because of the plethora of hnp information that they contain, do not collect consumption or income data; but they do have detailed information on households' physical characteristics, and on the household-level presence of and access to a wide range of goods and services. While there is some disagreement about the relative merits of using such wealth/asset information or consumption data to measure economic status, several recent studies suggest that the asset-consumption relationship is quite close.¹⁹ To the extent this is the case, an index of wealth or asset possession/availability can be taken as a

¹⁹ See, for example, Deon Filmer and Lant H. Pritchett, "Estimating Wealth Effects without Expenditure – or Tears: An Application to Educational Enrollments in States of India," *Demography* 38, no. 1 (February 2001): 115-32; Shea Rutstein and Kiersten Johnson, *The DHS Wealth Index*, DHS Comparative Reports No. 6 (Calverton, Maryland, USA: ORC Macro, August 2004) esp. 10-14; Adam Wagstaff and Naoko Watanabe, "What Difference Does the Choice of SES Make in Health Inequality Measurement?" *Health Economics* 12 (2003): 885-90.

reasonably satisfactory proxy for consumption, rather than or in addition to serving as an indicator of economic status in its own right.

C. ECONOMIC STATUS AS A MEASURE OF SOCIO-ECONOMIC WELL-BEING

Like consumption or income, a wealth index defines disparities that are primarily economic. This is by no means the only way to define inter-group inequalities that are of potential concern. Other possibilities include gender, place of residence, education, ethnic background, or other factors associated with social exclusion. Thus the economic perspective provides only a partial view of the multidimensional concepts of poverty, inequality, and inequity.

By including tables on female-male and rural-urban inequalities, this report pays adequate attention to two of inequality's other important dimensions to justify the use of the term "socio-economic" rather than simply "economic" in its title. However, the centrality of tabulations based on the wealth index means that the primary focus is on the economic dimension. The justification for this lies not in the greater importance of economic considerations, but rather in the recently-improved ability to analyze and thus begin dealing with them. Until the development of the wealth/asset approach, the assessment of economic status had been based on consumption, expenditures, or income, all of which are far more difficult to measure than such other, non-economic dimensions of inequality as gender, ethnic identity, educational level, and place of residence. As a result, assessments of health inequalities by economic status had lagged well behind measurements in terms of these other dimensions, especially gender and education. The focus on inequality's economic aspect applied here represents an effort to redress this imbalance.

D. INDEX CONSTRUCTION

Choice of Items

Use of a wealth index requires decisions about which items to include in it. In the case of secondary analyses like the one featured in this report, the choice is limited to those items included in the data sets being used. But even with this constraint, there nonetheless remains considerable room for choice, given the large number of items for which information is collected by the DHS.

The decision made in preparing this report was to include all items in each DHS household questionnaire that relate to ownership of household goods; to dwelling unit construction and characteristics; and to access to services and resources like electricity, water, and sanitation facilities. Also included were other potential indicators of wealth, such as live-in domestic servants. This decision, admittedly somewhat arbitrary, has both advantages and disadvantages.

The principal advantage is practical: use of a large number of assets increases the degree of variation across household asset scores and facilitates a more regular distribution of individuals across quintiles. It also reduces the possibility of subjectivity in selecting only some of the variables for inclusion on some *a priori* basis; and it may increase a wealth index's accuracy as a proxy for consumption.

However, including all variables is far from satisfying conceptually. For example, it means failing to discriminate with respect to the items' differing natures. It is not clear, for instance, whether access to water, sanitation, electricity, or other publicly-provided resources should be included in an index that purports to measure private household wealth.

Further, many items that are candidates for inclusion in a DHS-based wealth index might be seen as directly influencing health status: water and sanitation for infant and child mortality, for

example. It would be desirable to include quintile-specific estimates for such items; but to the extent that such items have large index coefficients, any estimates for those items would be suspect. Such items appear to be relatively few and of limited statistical significance in the index used here. However, for the sake of caution, quintile-specific estimates for items appearing in the index have nonetheless been excluded from the basic tables and appear only in supporting table III.C

Additional issues arise when comparing the findings for two different points in time covered in the basic tables. Because the nature and number of asset questions included in DHS surveys has been evolving, the items included in the wealth index differ somewhat for each of the surveys reported upon. As such, the results presented in the basic tables might differ to some extent from findings produced by some other approach, such as including in the index only those items appearing in each survey covered.

Weighting of Items

A further decision required in construction of an index concerns the weight to attach to each of the respective items. As noted earlier, the method used in this report is principal components analysis (PCA).

Adoption of this method was based on the findings, referred to earlier, that its use resulted in outcomes that approximated reasonably well those produced by taking a consumption or expenditure approach. Further, it often provides greater discrimination in economic status than does the use of consumption/expenditures. It has also emerged as the standard approach for use in analyses of the sort presented here, so that its adoption is largely non-controversial.

Yet this choice, too, is not without an arbitrary aspect; for alternative plausible methods exist. Examples include the “inverse possession” approach, which gives more weight to items possessed by only a few and less to those possessed by many;²⁰ or, perhaps, the common practice of simply assigning the same weight to each index item.

Also, the weights for any particular item vary from survey to survey, since the weights were determined separately for the population of each survey included in the basic tables. The results thus produced can be expected to differ from those generated in some other manner, such as generating common weights for all the surveys covered by pooling the data sets.

Use of Principal Components Analysis with Dichotomous Variables

An additional issue concerns the use of a technique like PCA, developed for use with continuous variables, in the construction of an index based primarily on dichotomous variables. While legitimate in principle, any reservations in this regard are of limited practical consequence, since the considerable experimentation undertaken in preparation for the tabulations presented here indicated that any inaccuracy introduced by applying PCA to the analysis of the dichotomous values used is minimal.

Economies of Scale

Calculating the values for a household wealth index also requires a decision concerning economies of scale that exist in the households covered. The calculations presented here assume complete economies of scale. The implicit assumption is that additional members do not add to household expenses on items included in the index.

²⁰ Saul S. Morris et al., “Validity of Rapid Estimates of Household Wealth and Income for Health Surveys in Rural Africa,” *Journal of Epidemiology and Community Health* 54 (2000): 381-87.

E. DEFINITION OF QUINTILES

Quintiles of Individuals

As noted earlier, the quintile-specific figures presented in this report refer to quintiles of individuals in the household population. Such quintiles need to be distinguished from quintiles of households or quintiles of only those people in the population who are “at risk”: that is, subject to the particular condition, eligible for the particular service, or capable of behaving in a particular way (children born alive for infant and under-five mortality, for example; or adult men and women for condom use or non-regular sexual partnerships).

The expression of findings in terms of quintiles of individuals has several implications:

- Because fertility is often higher in lower economic households than among better-off ones, the number of individuals per household will frequently be larger among the poor than among higher-income groups. In such cases, the number of households will vary systematically across quintiles of individuals, and the results expressed in terms of quintiles of households can differ significantly from those presented here.

- The proportion of individuals “at risk” with regard to a particular indicator is also likely to vary across quintiles in many cases. (For example, in cases where fertility is higher among poor people, a higher-than-average proportion of poor populations will consist of newborns at risk from infant mortality, young children subject to malnutrition, and pregnant women for whom antenatal care is relevant.) To facilitate the work of any investigators wishing to undertake calculations based on people at risk, the number of such people in each quintile of individuals is shown in part III.A.

- As previously indicated, the population average figure provided for each indicator is equivalent to the weighted sum of the quintile rates for that indicator, where the weight assigned to each quintile rate is the number of people at risk in each quintile as presented in part III.A. As a result of this weighting, the population average will usually differ from a simple mean of the population quintile estimates.

Quintiles of Males and Females, of Rural and Urban Residents

As also reported in the data and methods section, the tables on rural and urban residents and on men and women were prepared using the same asset scores as for the total population; and rural-urban residents and females-males were separated from one another only after the entire sample had been disaggregated into quintiles of individuals. This means that the figures given in the rural-urban and female-male tables refer to females-males and rural-urban residents belonging to each quintile of individuals in the total population, as distinct from quintiles of females, of males, of rural residents, or of urban residents alone.

The consequence of this distinction is particularly evident with regard to rural and urban residents. Since rural residents tend to be poorer than urban dwellers, they normally form a considerably higher proportion of individuals in the lower economic quintiles of the total population than in the higher ones. Conversely, urban residents tend to be concentrated in the higher economic groups. As a result, the number of individuals in each of the urban and rural quintiles usually varies greatly and systematically; and when this is the case, the figures presented in the rural-urban tables can differ significantly from those produced by a computation procedure that places the same number of rural and urban residents in each rural quintile or each urban quintile. (The results may also differ significantly from application of an approach featuring the separate calculation of index values for urban and for rural groups. While such separate index values may well be preferable conceptually, their calculation involves complexities that prevented their preparation for this report.)

F. COMPARISON OF QUINTILES ACROSS COUNTRIES

Reliance on population quintiles as basic presentational format for the data appearing in this report implicitly incorporates a relative concept of poverty. This differs from an absolute concept of poverty under which the population would be divided into groups of different sizes according to some absolute standard of living (such as people earning less than one dollar a day, between one and two dollars a day, and more than two dollars a day).

This means that, when comparing values of an indicator among people in a given quintile across countries, the comparison is between groups of people whose economic status can be quite different. The lowest quintile of a Latin American population, for example, will usually be considerably better-off than the lowest quintile in an African country.

G. COMPARISON OF QUINTILES OVER TIME

Another implication is that the wealth status of any given quintile within a particular country is likely to change over time. For instance, when a country is progressing economically, the wealth of the households in the population will tend to increase. This will raise the average asset score in most, possibly all population quintiles. As a result, the living standard enjoyed by individuals in any quintile covered by a recent survey is likely to be higher than that of individuals in that same quintile as measured in a prior survey.

H. STATISTICAL INDICATORS OF INEQUALITY

The available statistical indicators of inequality are far too numerous to permit use of more than a small proportion of them in presenting the findings featured in this report. The three indicators employed have been selected to provide a wide range of perspectives. Two are designed for ease of understanding, the third for greater technical accuracy.

The low/high quintile ratio and low-high quintile difference are the two presented for ease of understanding. The former is a relative measure, the latter an absolute measure that can produce a significantly different impression from that provided by the former.

The concentration index is provided for the benefit of technical specialists wishing greater accuracy. It measures the degree of inequality in an hnp indicator across the full wealth index distribution, rather than differences between only two of the five quintiles, and also reflects the relative size of the different asset-based divisions of the study population.

I. COMPARABILITY WITH OTHER REPORTS

Tabulations similar to those presented here can also be found in the initial series of hnp/poverty country reports, issued in 2000, and in the recent country reports issued by the DHS program. The figures presented in those reports normally resemble quite closely those appearing here for any given indicator; but there are often slight differences for one or more of several reasons. The most common is a difference in the definition of the indicator in question. (These differences are usually small and subtle. But there is one important exception: the definition of moderate malnutrition among children. In the 2000 reports, this was defined as second and third degree malnutrition taken together. Here, it is defined as second degree malnutrition alone.) Another reason, with respect to infant and child mortality, is that the figures in the DHS documents are typically based on experience during the five years before the survey in question, rather than

during the ten previous years as in this report. A further frequent reason is the use of an improved computational technique.

In addition, asset-based, quintile-specific tabulations of hnp indicators have begun to appear in an increasing number of other documents. Given the basic similarity of approach, such tabulations usually produce results that are generally congruent with those shown in part I of this report. However, significant divergences have occasionally been reported; and the absence of adequately detailed information about data and methods often prevents any fully-satisfying understanding of the approaches used. As a result, occasional doubts and frequent uncertainties about approach comparability remain.

PART IV. SUPPORTING TABLES, 2001

- A. SAMPLE SIZES**
- B. STANDARD ERRORS**
- C. ASSET DISTRIBUTION AND WEIGHTS**

Mali
2001 - SAMPLE SIZES

TOTAL SAMPLE

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Number of household members						
All	13,129	13,113	13,117	13,167	13,096	65,622
Urban	261	676	1,028	3,719	11,544	17,228
Rural	12,869	12,437	12,089	9,448	1,552	48,394
Female	6,785	6,670	6,782	6,682	6,636	33,556
Male	6,344	6,443	6,334	6,485	6,459	32,064

Part I: HNP STATUS

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Mortality rates						
All	5,714	5,626	5,450	5,301	4,186	26,289
Urban	116	262	412	1,420	3,632	5,842
Rural	5,599	5,365	5,039	3,887	556	20,447
Female	2,924	2,733	2,635	2,574	2,056	12,922
Male	2,790	2,897	2,817	2,731	2,133	13,367
Prevalence of fever, diarrhea, acute respiratory infection						
All	2,356	2,243	2,293	2,244	2,000	11,143
Urban	53	103	166	590	1,731	2,643
Rural	2,303	2,141	2,127	1,653	269	8,493
Female	1,208	1,104	1,134	1,066	983	5,495
Male	1,148	1,139	1,159	1,178	1,017	5,641
Total fertility rate						
All	6,994	6,848	7,048	7,176	8,206	36,296
Urban	177	389	562	2,084	7,367	10,579
Rural	6,818	6,462	6,490	5,103	844	25,717
Age-specific fertility rate 15-19						
All	1,294	1,300	1,462	1,522	2,425	8,003
Urban	43	89	105	500	2,232	2,969
Rural	1,251	1,211	1,357	1,022	194	5,035
Children's nutritional status						
All	1,995	1,909	1,954	1,901	1,643	9,408
Urban	40	88	153	500	1,424	2,205
Rural	1,955	1,821	1,801	1,401	219	7,197
Female	1,025	940	941	894	803	4,603
Male	971	969	1,013	1,007	840	4,800
Children's anemia status						
All	665	527	493	495	407	2,587
Urban	20	25	37	114	359	555
Rural	644	502	456	382	48	2,032
Female	318	273	238	285	203	1,317
Male	346	254	255	211	204	1,270

Mali
2001 - SAMPLE SIZES

Part I: HNP STATUS (Cont.)

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Women's nutritional status						
All	1,869	1,818	1,877	1,997	2,483	10,049
Urban	46	123	152	625	2,242	3,188
Rural	1,823	1,695	1,724	1,372	241	6,855
Women's anemia status						
All	895	655	641	767	827	3,787
Urban	25	38	43	228	747	1,081
Rural	870	617	598	540	80	2,705
Girls' circumcision						
All	1,852	1,812	1,824	1,800	1,561	8,855
Urban	45	92	148	503	1,369	2,161
Rural	1,806	1,720	1,675	1,297	191	6,694
Women's circumcision						
All	2,435	2,392	2,458	2,543	3,012	12,849
Urban	61	139	197	751	2,712	3,863
Rural	2,375	2,254	2,261	1,791	300	8,986
Prevalence of genital discharge, ulcer, sore						
Female	2,320	2,280	2,337	2,340	2,524	11,801
Urban Female	58	128	190	680	2,258	3,314
Rural Female	2,262	2,153	2,147	1,661	267	8,490
Male	655	441	472	494	774	2,836
Urban Male	9	32	35	165	702	943
Rural Male	646	409	438	329	72	1,894

Mali

2001 - SAMPLE SIZES

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Immunization coverage						
All	451	471	432	442	401	2,197
Urban	15	26	27	133	346	547
Rural	436	445	405	309	55	1,650
Female	230	224	215	214	192	1,075
Male	221	247	217	227	210	1,122
Treatment of fever						
All	695	603	644	562	481	2,988
Urban	17	39	42	122	414	634
Rural	678	564	602	440	68	2,352
Female	316	304	331	316	259	1,526
Male	379	299	313	246	222	1,459
Treatment of acute respiratory infection						
All	246	201	248	189	217	1,102
Urban	4	15	14	32	181	246
Rural	241	186	234	157	36	854
Female	139	106	127	96	98	566
Male	107	94	122	94	119	536
Treatment of diarrhea						
All	491	461	474	380	261	2,069
Urban	12	21	19	79	213	344
Rural	479	439	455	301	48	1,722
Female	236	221	242	153	128	980
Male	255	240	232	227	133	1,087
Antenatal and delivery care						
All	1,718	1,694	1,714	1,668	1,492	8,291
Urban	43	81	126	453	1,312	2,015
Rural	1,675	1,612	1,588	1,215	180	6,270
Contraceptive prevalence						
Female	2,238	2,199	2,217	2,138	1,923	10,723
Urban Female	57	119	176	583	1,684	2,619
Rural Female	2,181	2,079	2,042	1,555	239	8,096
Male	562	370	392	379	412	2,122
Urban Male	8	20	22	115	363	528
Rural Male	554	349	370	264	49	1,586
Contraceptive source						
Female	94	79	76	155	344	748
Urban Female	1	5	12	72	311	401
Rural Female	93	74	64	83	33	348
Male	562	370	392	379	412	2,122
Urban Male	8	20	22	115	363	528
Rural Male	554	349	370	264	49	1,586
Treatment of genital discharge, ulcer, sore						
Female	532	487	480	532	876	2,907
Urban Female	23	25	36	191	804	1,079
Rural Female	510	462	443	341	72	1,828
Male	44	36	30	30	45	185
Urban Male	1	6	3	10	42	62
Rural Male	43	31	27	20	3	124

Mali

2001 - SAMPLE SIZES

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Sanitary disposal of stools						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Wash hands prior to preparing food						
All	2,435	2,392	2,458	2,543	3,012	12,840
Urban	61	139	197	751	2,712	3,860
Rural	2,375	2,254	2,261	1,791	300	8,981
Handwashing facilities in household?						
All	2,315	2,379	2,747	2,649	2,229	12,319
Urban	46	124	219	711	1,959	3,059
Rural	2,269	2,255	2,528	1,938	271	9,261
Bednet ownership						
All	2,315	2,379	2,747	2,649	2,229	12,319
Urban	46	124	219	711	1,959	3,059
Rural	2,269	2,255	2,528	1,938	271	9,261
Bednet use by children						
All	1,588	1,571	1,626	1,590	1,357	7,732
Urban	32	80	110	422	1,179	1,823
Rural	1,556	1,491	1,516	1,168	178	5,909
Bednet use by pregnant women						
All	379	346	332	318	258	1,633
Urban	8	8	25	71	231	343
Rural	372	338	307	247	27	1,291
Exclusive breastfeeding						
All	201	202	209	190	159	961
Urban	7	4	9	45	145	210
Rural	193	198	200	145	14	750
Female	96	100	100	89	90	475
Male	105	102	109	100	69	485
Timely complementary breastfeeding						
All	166	184	196	195	174	915
Urban	0	8	18	51	144	221
Rural	166	175	178	144	30	693
Female	69	102	108	113	77	469
Male	97	81	89	82	97	446
Bottle-feeding						
All	561	520	597	572	498	2,748
Urban	14	17	40	134	428	633
Rural	547	503	556	438	70	2,114
Female	266	276	307	290	245	1,384
Male	295	244	289	282	253	1,363

Mali

2001 - SAMPLE SIZES

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR (Cont.)

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Iodized salt in household						
All	1,615	1,589	1,966	1,774	1,323	8,267
Urban	38	85	171	440	1,148	1,882
Rural	1,577	1,504	1,795	1,334	175	6,385
Vitamin A supplementation						
All	2,038	1,967	1,975	1,939	1,740	9,659
Urban	42	95	147	519	1,503	2,306
Rural	1,997	1,872	1,828	1,420	237	7,354
Female	988	1,003	1,001	1,037	877	4,906
Male	1,050	964	975	902	863	4,754
Tobacco and alcohol use, casual sexual partners, condom use for casual sex						
Female	2,435	2,392	2,458	2,543	3,012	12,840
Urban Female	61	139	197	751	2,712	3,860
Rural Female	2,375	2,254	2,261	1,791	300	8,981
Male	764	568	553	587	916	3,388
Urban Male	10	37	41	197	826	1,111
Rural Male	754	532	511	391	90	2,278
Domestic violence						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na

Mali

2001 - SAMPLE SIZES

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
School completion (Grade 5)						
Female	2,661	2,584	2,664	2,709	3,171	13,789
Urban female	65	149	211	777	2,847	4,049
Rural female	2,596	2,436	2,453	1,932	323	9,740
Male	2,110	2,172	2,102	2,272	3,218	11,874
Urban male	33	116	167	722	2,911	3,948
Rural male	2,077	2,057	1,935	1,550	308	7,926
School participation						
Female	1,274	1,229	1,148	1,145	944	5,741
Urban female	21	63	82	308	824	1,297
Rural female	1,252	1,167	1,067	837	121	4,443
Male	1,160	1,243	1,166	1,114	902	5,584
Urban male	16	46	80	324	753	1,219
Rural male	1,144	1,197	1,086	790	149	4,365
Mass media exposure						
Female	2,435	2,392	2,458	2,543	3,012	12,849
Urban female	61	139	197	751	2,712	3,863
Rural female	2,375	2,254	2,261	1,791	300	8,986
Male	764	568	553	587	916	3,405
Urban male	10	37	41	197	826	1,125
Rural male	754	532	511	391	90	2,280
Knowledge of HIV/AIDS prevention						
Female	4,851	2,392	2,458	2,543	3,012	15,256
Urban female	61	139	197	751	2,712	3,860
Rural female	2,375	2,254	2,261	1,791	300	8,981
Male	764	568	553	587	916	3,405
Urban male	10	37	41	197	826	1,111
Rural male	754	532	511	391	90	2,278
Household decisionmaking and justification of violence						
All	2,435	2,392	2,458	2,543	3,012	12,849
Urban	61	139	197	751	2,712	3,863
Rural	2,375	2,254	2,261	1,791	300	8,986
Orphanhood						
All	6,813	6,688	6,466	6,405	5,744	32,123
Urban	141	320	512	1,769	4,953	7,698
Rural	6,672	6,368	5,954	4,635	790	24,426
Female	3,451	3,325	3,231	3,144	2,998	16,153
Male	3,362	3,363	3,233	3,261	2,743	15,966

Mali

2001 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part I: HNP STATUS

Indicator	Wealth Quintiles					Avg.
	Low	2nd	3rd	4th	High	
A. Childhood illness and mortality						
Infant mortality rate	7.63	6.23	7.53	6.54	5.83	3.32
Under-five mortality rate	8.74	9.19	8.62	10.81	8.81	5.48
Prevalence of fever	1.48	1.61	1.21	1.37	1.48	0.78
Prevalence of diarrhea	1.18	1.33	1.13	0.89	1.01	0.61
Prevalence of acute respiratory infection	1.13	0.87	0.82	0.91	1.02	0.49
B. Fertility						
Total fertility rate	0.19	0.21	0.16	0.24	0.23	0.12
Adolescent fertility rate	13.00	14.00	11.00	14.00	10.00	8.00
C. Nutritional status						
<i>Children:</i>						
Moderate stunting	1.08	1.41	1.06	1.09	1.25	0.58
Severe stunting	1.30	1.59	1.22	1.09	1.17	0.68
Moderate underweight	1.04	1.14	1.12	1.18	1.31	0.62
Severe underweight	1.00	1.11	1.00	0.71	0.59	0.52
Mild anemia	1.78	2.86	1.79	1.95	1.98	0.95
Moderate anemia	2.75	3.81	2.85	2.90	3.26	1.75
Severe anemia	1.53	1.86	2.30	1.95	1.34	0.98
<i>Women:</i>						
Malnutrition	1.17	1.12	1.14	1.05	0.66	0.53
Mild anemia	2.11	2.17	2.09	2.57	2.09	0.93
Moderate anemia	1.60	1.88	1.80	2.25	1.50	0.80
Severe anemia	0.65	0.94	0.74	0.59	0.71	0.33
D. Female circumcision						
<i>Prevalence of circumcision:</i>						
Girls	2.10	1.87	2.26	2.02	1.89	1.32
Women	0.73	0.83	1.65	1.86	1.20	0.80
<i>Prevalence of occlusion:</i>						
Girls	0.39	0.47	0.58	0.56	0.81	0.29
Women	0.26	0.31	0.38	0.47	0.42	0.18
E. Sexually transmitted disease						
<i>Prevalence of genital discharge:</i>						
Women	1.39	1.61	1.77	1.54	1.50	0.90
Men	1.36	1.35	1.43	1.11	0.94	0.61
<i>Prevalence of genital ulcer:</i>						
Women	0.88	0.84	0.75	1.19	0.90	0.45
Men	0.89	1.01	0.44	0.54	0.44	0.32

Mali

2001 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization						
BCG coverage	3.82	3.26	3.26	2.75	1.73	1.76
Measles coverage	3.75	3.43	3.03	3.03	2.63	1.80
DPT coverage	3.16	3.16	2.95	3.98	3.43	2.14
Full basic coverage	3.14	2.81	2.51	3.26	3.68	1.85
No basic coverage	3.16	2.67	3.00	2.49	1.60	1.47
Hepatitis B coverage	na	na	na	na	na	na
Yellow fever coverage	na	na	na	na	na	na
B. Treatment of childhood illnesses						
<i>Treatment of fever:</i>						
Medical treatment of fever	2.40	2.25	2.31	2.98	2.81	1.35
Treatment in a public facility	1.14	1.32	1.78	1.92	3.20	0.99
Treatment in a private facility	2.03	1.81	1.84	2.08	2.83	1.00
<i>Treatment of acute respiratory infection (ARI):</i>						
Medical treatment of ARI	3.97	4.22	2.87	4.43	3.73	2.03
Treatment in a public facility	1.95	2.36	2.51	3.69	4.83	1.70
Treatment in a private facility	2.92	3.85	2.19	2.65	3.57	1.48
<i>Treatment of diarrhea:</i>						
Use of oral rehydration therapy	3.01	3.52	2.85	2.61	2.71	1.65
Medical treatment of diarrhea	1.79	1.59	2.01	2.17	3.88	1.17
Treatment in a public facility	0.85	0.79	1.10	1.93	3.49	0.81
Treatment in a private facility	1.65	1.50	1.65	1.40	3.73	0.85
C. Antenatal and delivery care						
<i>Antenatal care (ANC) visits:</i>						
To a medically trained person	2.70	2.68	2.42	2.34	0.93	1.85
To a doctor	0.22	0.34	0.47	0.43	1.21	0.30
To a nurse or trained midwife	2.42	2.52	2.32	2.24	1.43	1.52
Multiple visits to a medically trained person	1.80	2.67	2.26	2.24	1.92	1.47
<i>Antenatal care content:</i>						
Tetanus toxoid	2.22	2.15	1.90	2.37	1.20	1.37
Prophylactic antimalarial treatment	2.07	2.01	1.81	2.04	1.21	1.19
Iron supplementation	1.52	1.61	1.33	1.95	2.29	0.96
<i>Delivery attendance:</i>						
By a medically trained person	2.32	2.82	2.67	3.17	1.19	2.20
By a doctor	0.16	0.16	0.33	0.47	0.68	0.19
By a nurse or trained midwife	2.30	2.82	2.64	3.09	1.31	2.15
In a public facility	2.16	2.50	2.63	3.03	1.66	1.79
In a private facility	0.20	0.33	0.09	0.21	0.70	0.16
At home	2.18	2.46	2.59	3.11	1.43	1.83
D. Contraceptive services						
<i>Contraceptive prevalence:</i>						
Women	0.65	0.63	0.56	1.03	1.49	0.47
Men	1.93	1.65	1.69	2.18	2.54	0.99
<i>Source of contraception - public sector:</i>						
Women	5.99	7.26	7.37	5.10	3.12	2.32
Men	na	na	na	na	na	na

Mali

2001 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
D. Contraceptive services (cont.)						
<i>Source of contraception - private sector:</i>						
Women	3.42	6.33	4.56	3.81	2.93	1.95
Men	na	na	na	na	na	na
E. Treatment of adult illnesses						
<i>Treatment of genital discharge, ulcer, sore:</i>						
Women	1.86	2.42	2.45	2.47	2.35	1.75
Men	8.85	8.17	7.90	12.21	8.56	4.66
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>						
Women	1.39	1.61	1.77	1.54	1.50	0.90
Men	1.52	1.60	1.46	1.19	1.03	0.65
<i>Voluntary counseling and testing for HIV/AIDS:</i>						
Women	0.18	0.27	0.28	0.66	0.76	0.31
Men	1.03	1.35	0.93	1.52	1.99	0.76

Mali

2001 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles					Avg.
	Low	2nd	3rd	4th	High	
A. Hygienic practices						
<i>Disposal of children's stools:</i>						
Sanitary disposal	na	na	na	na	na	na
<i>Handwashing:</i>						
Wash hands prior to preparing food	0.92	0.98	1.09	0.91	0.73	0.51
Handwashing facilities in household	0.64	1.00	1.49	1.09	2.11	0.88
B. Bednet ownership and use						
<i>Bednet ownership:</i>						
Bednet ownership	2.46	2.10	2.02	2.18	2.56	1.43
Treated bednet ownership	na	na	na	na	na	na
<i>Bednet use:</i>						
By children	2.25	2.38	1.94	2.25	2.42	1.37
By pregnant women	3.66	3.52	3.95	3.59	4.46	1.96
C. Breastfeeding						
Exclusive breastfeeding	3.66	4.48	3.66	3.87	3.71	1.92
Timely complementary feeding	4.05	3.74	4.68	3.99	4.52	2.04
Bottle-feeding	0.58	0.87	0.52	1.02	1.93	0.51
D. Micronutrient consumption						
<i>Iodized salt:</i>						
Availability of iodized salt in household	1.88	1.92	1.67	1.93	2.28	1.06
<i>Vitamin A:</i>						
Children	2.33	2.06	2.24	3.01	2.50	1.28
Women	1.13	1.68	1.21	1.85	1.87	0.88
E. Tobacco and alcohol use						
<i>Tobacco:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na
<i>Alcohol:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na
F. Sexual practices						
<i>Non-regular sexual partnerships:</i>						
Women	0.21	0.13	0.17	0.27	0.44	0.13
Men	0.68	0.95	1.31	1.63	1.51	0.69
<i>Condom usage with non-regular partner:</i>						
Women	*	*	*	*	4.98	3.81
Men	11.33	9.53	10.49	7.44	4.94	3.58
G. Domestic violence						
Ever experienced violence	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na

Mali

2001 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
A. Education						
<i>School completion:</i>						
Women	0.40	0.44	0.51	1.13	2.07	0.93
Men	1.10	0.80	1.08	1.35	2.36	1.19
<i>School participation:</i>						
Girls	1.85	1.79	2.35	2.04	3.14	1.39
Boys	1.81	1.79	2.24	2.19	2.46	1.24
B. Exposure to mass media						
<i>Newspaper readership:</i>						
Women	0.12	0.17	0.25	0.48	1.28	0.47
Men	0.50	0.88	1.07	1.24	3.24	1.09
<i>Radio listenership:</i>						
Women	2.48	2.09	2.09	1.71	1.16	1.39
Men	2.29	2.45	2.10	1.77	1.35	1.06
<i>Television viewership:</i>						
Women	1.62	1.21	1.19	2.13	1.52	1.31
Men	2.78	3.02	2.66	3.28	2.31	2.02
C. Knowledge and attitudes about HIV/AIDS						
<i>Knowledge about sexual transmission of HIV/AIDS:</i>						
Women	2.19	2.42	2.07	1.88	1.10	1.49
Men	1.93	3.05	2.60	2.68	1.80	1.32
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>						
Women	1.71	2.04	2.24	1.68	1.19	1.24
Men	2.32	3.29	2.89	2.78	2.47	1.45
<i>Attitudes toward HIV/AIDS:</i>						
Women	1.71	2.36	1.45	1.57	1.18	0.96
Men	2.89	3.00	2.59	2.94	2.52	1.48
D. Status of women						
<i>Household decisionmaking:</i>						
Can seek own health care	0.79	0.84	1.03	1.28	0.80	0.52
Can seek children's health care	1.95	2.19	1.87	1.61	1.66	1.12
Can make daily household purchases	1.49	1.29	1.52	1.60	1.36	0.85
Can make large household purchases	1.13	1.19	1.28	1.24	1.30	0.70
Can make meal-related decisions	1.74	1.82	1.85	1.81	1.59	1.06
<i>Freedom of movement:</i>						
Can travel to visit family/relatives	1.22	1.29	1.04	2.43	1.56	0.86
<i>Other decisionmaking, attitudes:</i>						
Can decide how to spend own money	1.79	1.90	1.69	1.73	1.37	0.83
Can decide whether to have sex	2.07	1.97	1.96	2.04	1.70	1.23
Justifies domestic violence	1.14	0.91	1.27	0.98	1.38	0.68
E. Orphanhood						
Paternal orphan prevalence	0.41	0.47	0.44	0.50	0.72	0.25
Maternal orphan prevalence	0.32	0.36	0.31	0.28	0.31	0.14
Double orphan prevalence	0.13	0.27	0.16	0.14	0.15	0.09

Mali
2001 - ASSET DISTRIBUTION AND WEIGHTS
(FACTOR SCORE)

Asset Variable	Unweighted		Wealth Quintiles						Factor Score
	Mean	Std. Deviation	Low	2nd	3rd	4th	High	Avg.	
			Percentage of Population						
Has electricity	0.108	0.311	0.0%	0.7%	1.6%	5.4%	56.3%	12.8%	0.13708
Has radio	0.675	0.468	62.3%	73.1%	63.8%	76.5%	91.9%	73.5%	0.03933
Has television	0.129	0.335	0.0%	1.0%	4.1%	14.2%	69.8%	17.8%	0.12626
Has refrigerator	0.051	0.220	0.0%	0.0%	0.0%	0.1%	34.5%	6.9%	0.12706
Has bicycle	0.447	0.497	68.1%	67.0%	51.3%	46.7%	43.8%	55.4%	-0.02041
Has motorcycle/scooter	0.188	0.391	11.3%	25.1%	15.0%	26.9%	46.5%	24.9%	0.04393
Has car/truck	0.041	0.198	0.0%	0.1%	1.0%	3.5%	25.7%	6.1%	0.09137
Has telephone	0.023	0.151	0.0%	0.0%	0.0%	0.2%	17.1%	3.5%	0.09843
Has horse-drawn cart	0.391	0.488	92.3%	66.7%	26.0%	30.2%	11.9%	45.4%	-0.06888
Has plow	0.448	0.497	96.5%	79.3%	36.7%	31.6%	8.1%	50.4%	-0.07916
Has horse	0.090	0.287	23.1%	7.6%	3.9%	4.1%	1.2%	8.0%	-0.03329
Has wheel barrow	0.016	0.126	2.6%	1.1%	1.8%	1.2%	0.6%	1.4%	-0.00560
Has donkey	0.439	0.496	91.2%	71.3%	32.2%	33.1%	9.6%	47.5%	-0.07411
Has a domestic worker not related to head	0.012	0.107	0.0%	0.0%	0.0%	0.0%	11.5%	2.3%	0.06996
Works own or family's agric. land	0.275	0.447	59.4%	39.8%	38.2%	22.1%	3.2%	32.6%	-0.05072
Uses piped drinking water in residence	0.053	0.224	0.0%	0.0%	0.8%	2.2%	30.5%	6.7%	0.10174
Uses water that is piped into the building	0.023	0.150	0.0%	0.1%	1.7%	2.6%	7.3%	2.3%	0.03938
Uses a public faucet (piped)	0.185	0.388	6.1%	18.4%	15.5%	26.4%	35.2%	20.3%	0.03444
Uses an open well 1	0.093	0.291	5.3%	13.9%	8.5%	14.8%	10.7%	10.6%	-0.00341
Uses an open well 2	0.037	0.188	4.7%	4.5%	5.2%	5.1%	2.7%	4.4%	-0.00748
Uses an open well 3	0.375	0.484	68.3%	42.4%	40.6%	23.4%	4.3%	35.8%	-0.06403
Uses covered well 1 for drinking water	0.042	0.201	3.6%	4.8%	6.1%	5.9%	3.8%	4.8%	-0.00309
Uses covered well 2 for drinking water	0.018	0.131	1.0%	2.4%	2.2%	3.8%	2.3%	2.3%	0.00052
Uses covered well 3 for drinking water	0.088	0.283	8.3%	8.8%	9.5%	6.8%	2.0%	7.1%	-0.01824
Uses river, canal or surface water for drinking 2	0.068	0.252	1.3%	3.4%	8.0%	7.6%	0.9%	4.3%	-0.00871
Uses river, canal or surface water for drinking 3	0.006	0.079	0.8%	0.4%	0.7%	0.2%	0.0%	0.4%	-0.00609
Uses river, canal or surface water for drinking 4	0.000	0.009	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.00131
Uses spring for drinking water	0.008	0.087	0.2%	0.5%	0.9%	0.2%	0.0%	0.4%	-0.00527
Uses drinking water from tanker truck	0.000	0.013	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.00415
Uses bottled water	0.001	0.030	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.00142
Uses other source of drinking water	0.001	0.036	0.0%	0.1%	0.1%	0.3%	0.2%	0.1%	0.00137
Uses a flush toilet in residence/private	0.040	0.196	0.2%	1.7%	2.5%	2.8%	12.9%	4.0%	0.04780
Uses a pit latrine	0.292	0.455	36.8%	38.2%	35.5%	32.4%	22.2%	33.0%	-0.02121
Uses a ventilated improved pit latrine	0.039	0.194	0.2%	2.1%	2.6%	5.8%	12.0%	4.6%	0.02887
Uses a shared flush toilet in residence/private	0.025	0.156	0.2%	1.5%	1.5%	1.7%	5.1%	2.0%	0.02547
Uses a shared pit latrine	0.284	0.451	21.6%	27.2%	28.2%	36.2%	32.0%	29.0%	0.00984
Uses a shared ventilated improved pit latrine	0.056	0.230	0.4%	2.1%	4.4%	8.9%	15.4%	6.2%	0.03623
Uses bush,field as latrine	0.262	0.440	40.4%	27.2%	25.2%	11.3%	0.4%	20.9%	-0.05028
Uses other type of latrine	0.000	0.016	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.00062
Has dirt, sand, dung as principal floor in dwelling	0.837	0.370	100.0%	100.0%	99.4%	81.2%	15.3%	79.2%	-0.13250
Has wood, plank principal floor in dwelling	0.001	0.025	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.00224
Has tiles for main flooring material	0.000	0.020	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.00373
Has cement principal floor	0.002	0.043	0.0%	0.0%	0.2%	0.7%	0.1%	0.2%	0.00048
Has parquet or polished wood floors	0.013	0.112	0.0%	0.0%	0.0%	0.0%	9.7%	1.9%	0.07679
Has vinyl or asphalt strips as flooring material	0.143	0.350	0.0%	0.0%	0.1%	17.0%	73.4%	18.1%	0.11125
Has bricks for principal floor	0.003	0.053	0.0%	0.0%	0.0%	0.0%	1.3%	0.3%	0.02287
Has other type of flooring	0.000	0.016	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.00060

Mali
2001 - ASSET DISTRIBUTION AND WEIGHTS (Cont.)
(FACTOR SCORE)

Asset Variable	Unweighted		Wealth Quintiles						Factor Score
	Mean	Std. Deviation	Low	2nd	3rd	4th	High	Avg.	
			Percentage of Population						
Uses electricity as cooking fuel	0.000	0.016	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.00730
Uses gas as cooking fuel	0.005	0.070	0.0%	0.0%	0.0%	0.0%	1.3%	0.3%	0.03441
Uses biogas as cooking fuel	0.000	0.013	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.00447
Uses kerosene as cooking fuel	0.000	0.016	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.00183
Uses coal as cooking fuel	0.002	0.045	0.0%	0.0%	0.0%	0.1%	1.1%	0.2%	0.01636
Uses charcoal for cooking	0.100	0.299	0.0%	0.0%	1.6%	13.8%	34.4%	10.0%	0.09431
Uses wood as cooking fuel	0.825	0.380	99.8%	98.0%	91.9%	76.9%	62.0%	85.7%	-0.08159
Uses dung, manure as cooking fuel	0.054	0.226	0.1%	1.8%	6.2%	7.6%	0.5%	3.2%	-0.00525
Uses other cooking fuel	0.013	0.112	0.1%	0.2%	0.3%	1.0%	0.5%	0.4%	0.00447

PART V. SUPPORTING TABLES, 1995 / 96

- A. SAMPLE SIZES**
- B. STANDARD ERRORS**
- C. ASSET DISTRIBUTION AND WEIGHTS**

Mali
1995 / 96 - SAMPLE SIZES

TOTAL SAMPLE

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Number of household members						
All	9,735	9,773	9,767	9,735	9,762	48,772
Urban	531	823	1,321	3,318	7,991	13,983
Rural	9,204	8,950	8,446	6,417	1,772	34,789
Female	5,041	4,989	5,011	4,997	4,978	25,016
Male	4,694	4,784	4,756	4,738	4,784	23,756

Part I: HNP STATUS

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Mortality rates						
All	4,264	4,426	4,125	4,216	3,452	20,484
Urban	207	346	494	1,358	2,690	5,095
Rural	4,057	4,080	3,631	2,858	762	15,389
Female	2,140	2,226	2,041	2,086	1,723	10,217
Male	2,124	2,200	2,084	2,130	1,729	10,266
Prevalence of fever, diarrhea, acute respiratory infection						
All	1,029	1,096	1,072	1,100	940	5,237
Urban	44	85	131	381	759	1,400
Rural	985	1,011	941	719	181	3,837
Female	555	541	550	546	471	2,663
Male	474	555	522	553	469	2,573
Total fertility rate						
All	5,337	5,406	5,171	5,345	6,145	27,404
Urban	294	464	714	1,897	5,153	8,522
Rural	5,043	4,942	4,457	3,448	992	18,882
Age-specific fertility rate 15-19						
All	938	1,036	1,020	1,123	1,561	5,678
Urban	63	119	165	430	1,378	2,155
Rural	875	917	854	693	183	3,522
Children's nutritional status						
All	933	963	964	990	828	4,678
Urban	37	74	124	337	664	1,236
Rural	896	889	840	654	164	3,443
Female	509	490	495	494	402	2,390
Male	424	473	469	497	426	2,289
Children's anemia status						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Female	na	na	na	na	na	na
Male	na	na	na	na	na	na

Mali
1995 / 96 - SAMPLE SIZES

Part I: HNP STATUS (Cont.)

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Women's nutritional status						
All	810	855	820	828	713	4,026
Urban	44	67	99	287	572	1,069
Rural	766	788	721	541	141	2,957
Women's anemia status						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Girls' circumcision						
All	1,276	1,280	1,242	1,262	1,209	6,268
Urban	62	103	155	419	987	1,726
Rural	1,214	1,177	1,087	843	222	4,542
Women's circumcision						
All	1,870	1,900	1,823	1,891	2,221	9,704
Urban	103	168	257	679	1,867	3,074
Rural	1,767	1,732	1,565	1,213	354	6,630
Prevalence of genital discharge, ulcer, sore						
Female	1,774	1,791	1,721	1,767	1,951	9,004
Urban Female	97	149	234	619	1,624	2,723
Rural Female	1,677	1,642	1,487	1,147	328	6,281
Male	414	416	361	421	497	2,109
Urban Male	30	25	56	168	426	705
Rural Male	385	391	305	254	71	1,406

Mali

1995 / 96 - SAMPLE SIZES

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Immunization coverage						
All	290	332	313	349	277	1,562
Urban	13	24	31	120	226	414
Rural	277	308	283	229	51	1,148
Female	160	158	175	179	139	811
Male	130	174	138	170	139	751
Treatment of fever						
All	431	454	431	426	285	2,027
Urban	29	42	48	135	216	470
Rural	402	412	382	292	69	1,557
Female	212	220	222	216	160	1,030
Male	219	234	208	210	125	996
Treatment of acute respiratory infection						
All	159	182	161	168	130	800
Urban	7	11	16	54	105	193
Rural	151	171	144	114	25	605
Female	90	91	79	84	60	404
Male	69	91	82	85	70	397
Treatment of diarrhea						
All	298	294	298	283	153	1,326
Urban	17	23	30	85	113	268
Rural	281	271	268	198	40	1,058
Female	152	146	146	148	63	655
Male	147	148	152	135	89	671
Antenatal and delivery care						
All	1,055	1,102	1,040	1,078	901	5,175
Urban	49	85	129	359	723	1,345
Rural	1,006	1,016	911	719	178	3,830
Contraceptive prevalence						
Female	1,721	1,722	1,637	1,611	1,531	8,222
Urban Female	90	131	206	530	1,222	2,179
Rural Female	1,631	1,591	1,431	1,082	309	6,044
Male	370	372	288	321	294	1,645
Urban Male	22	24	42	127	239	454
Rural Male	349	348	246	194	56	1,193
Contraceptive source						
Female	8	25	33	68	235	369
Urban Female	1	0	12	32	207	252
Rural Female	7	24	21	36	27	115
Male	370	372	288	321	294	1,645
Urban Male	22	24	42	127	239	454
Rural Male	349	348	246	194	56	1,193
Treatment of genital discharge, ulcer, sore						
Female	na	na	na	na	na	na
Urban Female	na	na	na	na	na	na
Rural Female	na	na	na	na	na	na
Male	na	na	na	na	na	na
Urban Male	na	na	na	na	na	na
Rural Male	na	na	na	na	na	na

Mali

1995 / 96 - SAMPLE SIZES

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Sanitary disposal of stools						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Wash hands prior to preparing food						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Handwashing facilities in household?						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Bednet ownership						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Bednet use by children						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Bednet use by pregnant women						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Exclusive breastfeeding						
All	143	133	134	127	120	657
Urban	7	12	17	46	96	178
Rural	136	122	117	81	24	480
Female	63	69	64	59	56	311
Male	80	64	70	68	65	347
Timely complementary breastfeeding						
All	129	146	150	124	109	658
Urban	6	13	18	44	84	165
Rural	124	133	132	80	25	494
Female	52	76	77	65	55	325
Male	77	71	73	58	55	334
Bottle-feeding						
All	421	434	445	387	353	2,040
Urban	21	36	55	139	276	527
Rural	400	397	390	248	77	1,512
Female	181	221	220	197	175	994
Male	239	213	225	190	178	1,045

Mali

1995 / 96 - SAMPLE SIZES

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR (Cont.)

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
Iodized salt in household						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Vitamin A supplementation						
All	804	880	853	893	755	4,185
Urban	31	65	101	301	612	1,110
Rural	773	815	751	592	144	3,075
Female	378	448	414	449	379	2,068
Male	426	432	439	444	376	2,117
Tobacco and alcohol use, casual sexual partners, condom use for casual sex						
Female	1,870	1,900	1,823	1,891	2,221	9,705
Urban Female	103	168	257	679	1,867	3,074
Rural Female	1,767	1,732	1,565	1,213	354	6,631
Male	472	485	447	491	579	2,474
Urban Male	36	32	64	187	494	813
Rural Male	436	453	383	304	85	1,661
Domestic violence						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na

Mali

1995 / 96 - SAMPLE SIZES

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles					Pop. Total
	Low	2nd	3rd	4th	High	
School completion (Grade 5)						
Female	2,022	2,034	1,995	2,023	2,231	10,303
Urban female	113	168	273	702	1,858	3,114
Rural female	1,908	1,865	1,722	1,321	373	7,189
Male	1,615	1,653	1,628	1,729	2,140	8,766
Urban male	89	128	241	655	1,845	2,957
Rural male	1,526	1,525	1,387	1,074	296	5,809
School participation						
Female	886	885	837	838	816	4,262
Urban female	40	85	105	261	664	1,155
Rural female	846	800	732	577	152	3,106
Male	824	865	869	892	739	4,190
Urban male	48	85	110	281	573	1,097
Rural male	776	781	759	611	167	3,093
Mass media exposure						
Female	1,870	1,900	1,823	1,891	2,221	9,704
Urban female	103	168	257	679	1,867	3,074
Rural female	1,767	1,732	1,565	1,213	354	6,630
Male	472	485	447	491	579	2,474
Urban male	36	32	64	187	494	813
Rural male	436	453	383	304	85	1,661
Knowledge of HIV/AIDS prevention						
Female	3,723	1,900	1,823	1,891	2,221	11,558
Urban female	103	168	257	679	1,867	3,074
Rural female	1,767	1,732	1,565	1,213	354	6,631
Male	472	485	447	491	579	2,474
Urban male	36	32	64	187	494	813
Rural male	436	453	383	304	85	1,661
Household decisionmaking and justification of violence						
All	na	na	na	na	na	na
Urban	na	na	na	na	na	na
Rural	na	na	na	na	na	na
Orphanhood						
All	4,601	4,757	4,788	4,816	4,549	23,512
Urban	243	407	602	1,624	3,609	6,484
Rural	4,358	4,351	4,186	3,192	941	17,028
Female	2,718	2,805	2,595	2,709	2,419	13,246
Male	1,883	1,953	2,193	2,107	2,127	10,262

Mali

1995 / 96 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part I: HNP STATUS

Indicator	Wealth Quintiles					Avg.
	Low	2nd	3rd	4th	High	
A. Childhood illness and mortality						
Infant mortality rate	8.73	9.25	9.61	10.23	11.06	4.58
Under-five mortality rate	9.05	9.65	10.58	11.20	11.14	5.06
Prevalence of fever	1.73	1.86	1.82	1.99	2.18	0.98
Prevalence of diarrhea	1.66	1.39	1.54	1.42	1.45	0.75
Prevalence of acute respiratory infection	1.37	1.34	1.39	1.38	1.18	0.68
B. Fertility						
Total fertility rate	0.32	0.27	0.27	0.22	0.14	0.13
Adolescent fertility rate	20.00	20.00	20.00	10.00	10.00	10.00
C. Nutritional status						
<i>Children:</i>						
Moderate stunting	1.16	1.33	1.23	1.29	1.15	0.57
Severe stunting	1.52	1.34	1.20	1.26	0.99	0.66
Moderate underweight	1.47	1.54	1.39	1.42	1.64	0.73
Severe underweight	1.66	1.41	1.32	1.34	0.96	0.72
Mild anemia	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na
<i>Women:</i>						
Malnutrition	1.44	1.52	1.46	1.37	1.41	0.75
Mild anemia	na	na	na	na	na	na
Moderate anemia	na	na	na	na	na	na
Severe anemia	na	na	na	na	na	na
D. Female circumcision						
<i>Prevalence of circumcision:</i>						
Girls	3.47	1.87	1.55	1.47	1.34	1.15
Women	3.19	0.66	0.64	0.68	0.80	0.84
<i>Prevalence of occlusion:</i>						
Girls	na	na	na	na	na	na
Women	0.10	0.27	0.15	0.19	0.17	0.10
E. Sexually transmitted disease						
<i>Prevalence of genital discharge:</i>						
Women	0.09	0.15	0.12	0.17	0.12	0.06
Men	0.73	0.61	0.83	0.76	0.73	0.35
<i>Prevalence of genital ulcer:</i>						
Women	0.10	0.11	0.11	0.09	0.12	0.05
Men	0.69	0.99	0.40	1.02	0.42	0.36

Mali

1995 / 96 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
A. Childhood immunization						
BCG coverage	4.10	3.35	2.93	2.32	1.57	1.80
Measles coverage	3.75	3.27	3.51	3.34	2.71	1.92
DPT coverage	3.05	3.34	3.28	3.18	3.28	1.99
Full basic coverage	2.98	3.09	3.27	3.21	3.06	1.90
No basic coverage	4.13	3.35	2.92	2.15	1.51	1.80
Hepatitis B coverage	na	na	na	na	na	na
Yellow fever coverage	0.34	0.60	0.92	1.87	3.01	0.76
B. Treatment of childhood illnesses						
<i>Treatment of fever:</i>						
Medical treatment of fever	1.36	1.49	1.90	1.66	3.03	1.00
Treatment in a public facility	1.13	1.20	1.26	1.46	2.25	0.73
Treatment in a private facility	0.85	1.08	1.36	1.14	2.36	0.65
<i>Treatment of acute respiratory infection (ARI):</i>						
Medical treatment of ARI	3.71	2.97	3.60	3.84	4.36	2.26
Treatment in a public facility	3.10	2.75	2.69	3.43	5.02	1.84
Treatment in a private facility	2.05	1.96	2.93	3.01	3.78	1.38
<i>Treatment of diarrhea:</i>						
Use of oral rehydration therapy	3.16	2.85	3.35	3.19	4.68	1.76
Medical treatment of diarrhea	1.91	2.67	2.56	2.51	4.03	1.37
Treatment in a public facility	1.23	2.44	2.08	2.01	4.03	1.07
Treatment in a private facility	1.46	1.52	1.64	1.79	3.60	0.96
C. Antenatal and delivery care						
<i>Antenatal care (ANC) visits:</i>						
To a medically trained person	2.13	2.48	2.46	2.41	1.57	1.57
To a doctor	0.13	0.26	0.41	0.32	0.76	0.19
To a nurse or trained midwife	2.24	2.51	2.58	2.62	1.57	1.67
Multiple visits to a medically trained person	1.45	2.00	2.25	2.50	1.74	1.46
<i>Antenatal care content:</i>						
Tetanus toxoid	1.84	1.94	1.99	2.05	1.51	1.18
Prophylactic antimalarial treatment	na	na	na	na	na	na
Iron supplementation	na	na	na	na	na	na
<i>Delivery attendance:</i>						
By a medically trained person	1.34	2.14	2.79	2.78	1.90	1.80
By a doctor	0.08	0.25	0.18	0.43	0.62	0.17
By a nurse or trained midwife	1.33	2.15	2.77	2.75	1.89	1.77
In a public facility	1.20	1.68	2.00	2.83	2.36	1.51
In a private facility	0.09	0.00	0.18	0.14	0.54	0.11
At home	1.36	2.11	2.95	2.92	2.02	1.83
D. Contraceptive services						
<i>Contraceptive prevalence:</i>						
Women	0.18	0.44	0.51	0.76	1.31	0.38
Men	0.47	1.03	1.51	2.44	2.85	0.85
<i>Source of contraception - public sector:</i>						
Women	*	*	9.13	6.31	3.49	2.97
Men	na	na	na	na	na	na

Mali

1995 / 96 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part II: Intermediate Determinants of HNP Status - HNP SERVICE USE (Cont.)

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
D. Contraceptive services (cont.)						
<i>Source of contraception - private sector:</i>						
Women	*	*	9.15	6.25	3.63	3.02
Men	na	na	na	na	na	na
E. Treatment of adult illnesses						
<i>Treatment of genital discharge, ulcer, sore:</i>						
Women	*	*	*	*	11.28	6.30
Men	*	*	*	*	*	5.81
<i>Treatment of genital discharge, ulcer, sore in public facilities:</i>						
Women	0.09	0.15	0.12	0.17	0.12	0.06
Men	0.77	1.07	0.82	1.27	0.83	0.48
<i>Voluntary counseling and testing for HIV/AIDS:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na

Mali

1995 / 96 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part III: Intermediate Determinants of HNP Status - INDIVIDUAL AND HOUSEHOLD BEHAVIOR

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
A. Hygienic practices						
<i>Disposal of children's stools:</i>						
Sanitary disposal	na	na	na	na	na	na
<i>Handwashing:</i>						
Wash hands prior to preparing food	na	na	na	na	na	na
Handwashing facilities in household	na	na	na	na	na	na
B. Bednet ownership and use						
<i>Bednet ownership:</i>						
Bednet ownership	na	na	na	na	na	na
Treated bednet ownership	na	na	na	na	na	na
<i>Bednet use:</i>						
By children	na	na	na	na	na	na
By pregnant women	na	na	na	na	na	na
C. Breastfeeding						
Exclusive breastfeeding	3.31	3.37	3.36	3.58	3.05	1.56
Timely complementary feeding	4.74	3.66	3.88	4.26	4.43	1.87
Bottle-feeding	0.78	0.63	0.67	1.35	1.89	0.48
D. Micronutrient consumption						
<i>Iodized salt:</i>						
Availability of iodized salt in household	na	na	na	na	na	na
<i>Vitamin A:</i>						
Children	0.79	0.26	0.39	0.42	0.34	0.25
Women	na	na	na	na	na	na
E. Tobacco and alcohol use						
<i>Tobacco:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na
<i>Alcohol:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na
F. Sexual practices						
<i>Non-regular sexual partnerships:</i>						
Women	na	na	na	na	na	na
Men	1.29	1.55	1.83	1.94	2.79	1.02
<i>Condom usage with non-regular partner:</i>						
Women	na	na	na	na	na	na
Men	3.80	4.65	4.36	5.35	3.47	2.62
G. Domestic violence						
Ever experienced violence	na	na	na	na	na	na
Experienced violence in past year	na	na	na	na	na	na

Mali

1995 / 96 - STANDARD ERRORS OF QUINTILE ESTIMATES FOR TOTAL POPULATION

Part IV: UNDERLYING DETERMINANTS OF HNP STATUS

Indicator	Wealth Quintiles					
	Low	2nd	3rd	4th	High	Avg.
A. Education						
<i>School completion:</i>						
Women	0.27	0.38	0.56	0.87	1.77	0.69
Men	0.67	0.86	1.10	1.45	1.99	1.06
<i>School participation:</i>						
Girls	1.14	1.66	1.95	2.54	2.06	1.38
Boys	1.76	1.82	1.83	2.50	2.19	1.38
B. Exposure to mass media						
<i>Newspaper readership:</i>						
Women	0.15	0.24	0.28	0.59	1.43	0.51
Men	0.87	0.99	1.25	1.90	2.84	1.07
<i>Radio listenership:</i>						
Women	1.74	1.77	1.83	1.58	1.09	1.20
Men	2.51	2.25	2.00	1.42	1.11	1.11
<i>Television viewership:</i>						
Women	1.08	1.33	1.44	1.79	1.71	1.32
Men	1.88	2.69	3.07	3.47	2.51	1.97
C. Knowledge and attitudes about HIV/AIDS						
<i>Knowledge about sexual transmission of HIV/AIDS:</i>						
Women	2.38	1.78	1.69	1.26	0.73	1.05
Men	1.71	1.52	1.16	1.41	0.40	0.63
<i>Knowledge about mother-to-child transmission of HIV/AIDS:</i>						
Women	1.31	1.38	1.58	1.70	1.41	1.11
Men	2.83	2.43	2.85	2.53	2.15	1.34
<i>Attitudes toward HIV/AIDS:</i>						
Women	na	na	na	na	na	na
Men	na	na	na	na	na	na
D. Status of women						
<i>Household decisionmaking:</i>						
Can seek own health care	na	na	na	na	na	na
Can seek children's health care	na	na	na	na	na	na
Can make daily household purchases	na	na	na	na	na	na
Can make large household purchases	na	na	na	na	na	na
Can make meal-related decisions	na	na	na	na	na	na
<i>Freedom of movement:</i>						
Can travel to visit family/relatives	na	na	na	na	na	na
<i>Other decisionmaking, attitudes:</i>						
Can decide how to spend own money	2.97	2.40	2.37	1.68	1.16	1.28
Can decide whether to have sex	na	na	na	na	na	na
Justifies domestic violence	na	na	na	na	na	na
E. Orphanhood						
Paternal orphan prevalence	0.50	0.48	0.49	0.51	0.52	0.24
Maternal orphan prevalence	0.37	0.30	0.30	0.33	0.27	0.14
Double orphan prevalence	0.14	0.09	0.09	0.11	0.16	0.06

Mali

1995 / 96 - ASSET DISTRIBUTION AND WEIGHTS

(FACTOR SCORE)

Asset Variable	Unweighted		Wealth Quintiles						Factor Score
	Mean	Std. Deviation	Low	2nd	3rd	4th	High	Avg.	
			Percentage of Population						
Has electricity	0.070	0.256	0.0%	0.0%	0.0%	0.4%	37.5%	7.6%	0.17385
Has radio	0.568	0.495	21.4%	44.9%	69.0%	83.3%	92.2%	62.2%	0.08863
Has television	0.078	0.269	0.0%	0.0%	0.1%	1.4%	54.3%	11.2%	0.16504
Has refrigerator	0.033	0.178	0.0%	0.0%	0.0%	0.0%	20.7%	4.2%	0.16000
Has bicycle	0.351	0.476	36.5%	43.9%	53.3%	48.1%	29.5%	42.3%	-0.01605
Has motorcycle	0.163	0.369	0.8%	6.0%	12.3%	32.8%	54.5%	21.3%	0.09232
Has car	0.029	0.166	0.0%	0.0%	0.0%	0.0%	20.4%	4.1%	0.13180
Has telephone	0.007	0.083	0.0%	0.0%	0.0%	0.0%	5.3%	1.1%	0.10365
Uses piped drinking water in residence	0.057	0.232	0.0%	0.0%	0.0%	2.1%	26.2%	5.7%	0.13984
Uses piped drinking water in public tap	0.148	0.355	0.6%	2.1%	4.3%	14.7%	34.0%	11.1%	0.04957
Uses inside well drinking water	0.172	0.377	2.4%	8.6%	24.5%	38.3%	24.0%	19.6%	0.02008
Uses river, canal or surface water for drinking	0.040	0.197	7.6%	4.0%	4.7%	4.2%	0.9%	4.3%	-0.01777
Uses public well for water	0.438	0.496	77.6%	67.9%	43.7%	23.9%	8.0%	44.2%	-0.09247
Uses bore hole for water	0.142	0.349	11.7%	17.2%	21.6%	16.9%	6.7%	14.8%	-0.02349
Uses rain for drinking water	0.000	0.011	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.00147
Uses private flush toilet	0.004	0.062	0.0%	0.0%	0.0%	0.0%	2.5%	0.5%	0.06826
Uses a latrine	0.068	0.253	0.0%	1.6%	4.0%	10.8%	22.5%	7.8%	0.07023
Uses shared flush toilet	0.005	0.071	0.0%	0.0%	0.0%	0.0%	3.1%	0.6%	0.07293
Uses traditional pit latrine	0.620	0.485	9.9%	67.9%	85.0%	85.8%	70.9%	63.9%	0.03587
Uses bush, field as latrine	0.298	0.457	89.8%	29.9%	9.7%	3.2%	1.0%	26.7%	-0.09692
Uses other type of latrine	0.001	0.030	0.2%	0.2%	0.1%	0.0%	0.0%	0.1%	-0.00377
Has dirt, earth principal floor in dwelling	0.589	0.492	87.5%	73.2%	61.0%	40.7%	9.5%	54.4%	-0.10815
Has parquet or tile principal floor in dwelling	0.000	0.011	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.00048
Has cement principal floor	0.188	0.391	0.0%	0.0%	0.6%	24.6%	79.2%	20.9%	0.13350
Has tile flooring	0.010	0.098	0.0%	0.0%	0.0%	0.0%	5.4%	1.1%	0.10646
Has other type of flooring	0.000	0.011	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-0.00088
Has a dung floor	0.205	0.404	12.4%	26.6%	37.2%	34.2%	3.9%	22.9%	-0.02812
Has palm or bamboo flooring	0.001	0.030	0.1%	0.1%	0.0%	0.2%	0.0%	0.1%	-0.00212
Has vinyl/asphalt flooring	0.001	0.036	0.0%	0.0%	0.0%	0.0%	0.4%	0.1%	0.02270
Has carpeted flooring	0.002	0.047	0.0%	0.0%	0.0%	0.0%	1.0%	0.2%	0.02477
Rooms for sleeping	2.422	1.519	2.0	2.5	3.2	3.5	4.0	3.1	0.06486
Number of members per sleeping room	2.395	1.486	3.0	2.6	2.4	2.5	2.4	2.6	-0.01969

PART VI. ANNEXES

- A. SOURCES OF ADDITIONAL INFORMATION**
- B. USE OF INFORMATION FROM THIS REPORT TO MONITOR THE ECONOMIC STATUS OF PEOPLE SERVED BY HNP PROGRAMS**
- C. COUNTRIES COVERED BY THE HNP-POVERTY REPORT PROJECT**

ANNEX A. SOURCES OF ADDITIONAL INFORMATION

ADDITIONAL INEQUALITY DATA

World Bank HNP/Poverty Country Report Website:
<http://www.worldbank.org/hnp/povertyandhealth/countrydata>.

This World Bank website provides the full texts and tables for all fifty-six countries covered by the HNP/Poverty Country Report Project. (A list of the countries covered appears in annex C, at the end of this report.) Also available at the site are summary tables, organized by indicator, designed to facilitate cross-country comparisons in inequality with respect to particular indicators.

DHS Country Reports: <http://www.measuredhs.com/countries>.

All DHS final country reports produced since 2003 include quintile-specific tabulations in approximately 50-100 of the reports' HNP indicator tables. The tables deal with some of the indicators covered in this volume, and with many that are not.

UNICEF Multi-Indicator Cluster Survey Website:
<http://www.childinfo.org/MICS2/natlMICSrepz/MICSnatrep.htm>.

The UNICEF Multi-Indicator Cluster Survey (MICS) project is generally similar to the DHS program, but covers a somewhat different set of countries and indicators. The "standard tables" section for each country listed at the MICS website provides wealth-based, quintile-specific information in around 40-45 of the tables related to hnp, education, and child labor. In deriving these quintile-specific estimates, the MICS investigators have employed a wealth index similar to the one used here.

World Health Organization World Health Survey Website:
<http://www.who.int/healthinfo/survey/en/index.html>.

The World Health Organization's World Health Survey (WHS) includes such issues as self-assessed adult health status; coverage of interventions against adult chronic diseases and against maternal and child health problems; household health expenditures; insurance coverage; and health system responsiveness. Approximately seventy countries – developed as well as developing – have been covered thus far. Household wealth information has been collected and used to prepare quintile-specific estimates for many of the indicators appearing in the reports on these countries.

METHODS AND RESOURCES FOR FURTHER INEQUALITY ANALYSIS

Shea Oscar Rutstein and Kiersten Johnson, *The DHS Wealth Index*, DHS Comparative Reports No. 6 (Calverton, Maryland, USA: ORC Macro, August 2004) (Available at: http://www.measuredhs.com/pubs/pub_details.cfm?ID=470&srchTp=type).

This DHS publication, by two of the co-authors of the current report, describes in detail the construction of the wealth index that underlies the data presented in the basic tables.

Deon Filmer and Lant H. Pritchett, "Estimating Wealth Effects without Expenditure Data – or Tears: An Application to Education Enrollments in States in India," *Demography* 38, no.1 (February 2001): 115-132.

This seminal piece gave birth to the wealth index procedure used in the current volume. It also includes three of the previously-cited country case studies demonstrating the close relationship between results produced using wealth and those based on consumption as an indicator of household economic status.

Owen O'Donnell, Eddy van Doorslaer, Adam Wagstaff, and Magnus Lindelow. *Quantitative Techniques for Health Equity Analysis*. Washington D.C.: The World Bank, forthcoming.

Among the topics covered in this comprehensive overview of available quantitative techniques are the measurement of living standards using a wealth index and other approaches (chapter 6) and the concentration index as a measure of inequality (chapter 8).

DHS Country Data Sets: <http://www.measuredhs.com/accesssurveys/search>.

The data sets for all DHS surveys undertaken since 2003 include two pieces of information for each household that are designed to help investigators prepare quintile-specific tabulations for any indicator. These are: 1) the household wealth score; and 2) the economic quintile to which individuals in the household belong. Any tabulation using these pieces of information will be comparable to the figures appearing here.

ANNEX B. USE OF INFORMATION FROM THIS REPORT TO MONITOR THE ECONOMIC STATUS OF PEOPLE SERVED BY HNP PROGRAMS

The wealth or asset approach employed in this report can be used to monitor the economic status of people served by health, nutrition, and population (hnp) programs in two ways. The first, simpler way is suitable for monitoring nationwide, facility-based programs. A second, fuller version can also be employed for other types of programs, such as initiatives undertaken only in some parts of a country, or activities like mass education or outreach programs that do not operate through facilities.

BASIC MONITORING OF NATIONWIDE FACILITY-BASED PROGRAMS

The economic distribution of patients in a nationwide, facility-based program (say, a network of rural health posts, antenatal care clinics, emergency obstetrical facilities, or hospitals) can be determined through an exit survey of facility patients, using the wealth questionnaire and the set of quintile cut-off points that immediately follow this text, and which have been created using the information presented in part III.C. The questionnaire can be employed to measure the economic status of any individual responding to the questions on it. The set of cut-off points can serve to compare the distribution of the respondents' economic status with that of the nationally-representative sample of people interviewed by the DHS survey on which the present report is based.

The first step is to use the questionnaire in interviewing an adequately-large sample of patients attending the facility-based services of interest. The wealth score for each patient can then be calculated by multiplying the response to each question by the item scores also provided on the questionnaire, and summing the results. After this has been done, the quintile cut-off points can be used to place each individual in the economic quintile to which (s)he belongs. The number of patients and percentage of total patients in each quintile can then be calculated.

Since each quintile defined by the cut-off points contains 20 percent of the individuals in the nationally-representative DHS sample, the patients belonging to any such quintile containing significantly more (or less) than 20 percent of the total are over- (under-) represented relative to the national population. When the percentage of patients in each of the five quintiles is viewed as a whole, the result is a frequency distribution that indicates the spread of service beneficiaries across economic classes of individuals.²¹ For example:

- A service that favors the poorest people would have substantially more than 20 percent of its patients in each of the lowest one or two economic quintiles; considerably less than 20 percent of its patients in each of the highest quintiles.

²¹ That is, across economic classes of all individuals in the sample population. Estimates pertaining to quintiles of only those individuals needing services require adjusting the results of the procedure described here through application of the relevant quintile-specific, sample-size figures presented in part III.A.

- A service that reaches all economic classes equally would draw roughly the same proportion of total patients from the lower and upper quintiles.
- A service that favors the least poor population groups would have well over 20 percent of its patients in each of the highest one or two quintiles, considerably less than 20 percent of its patients in each of the lowest quintiles.

FULLER MONITORING OF FACILITY-BASED AND OF OTHER PROGRAMS

While capable of providing far more information than currently exists about the distribution of a program's beneficiaries, the approach just described has important limitations. For example, it cannot deal with the many important types of health programs that do not deliver services primarily through facilities – mass media health education, household visits by health workers, and many social marketing initiatives, for example. It is also limited in its ability to assess programs working only with certain areas within a country: it can compare the economic status of the programs' beneficiaries with that of the national population, but not with that of the specific sub-national areas where the programs are active. Further, it focuses primarily on only one of the two important dimensions of monitoring the distribution of program beneficiaries: that is, incidence or focus – the percentage of program benefits that flow to the poor. It cannot deal nearly so well with the second dimension, which concerns coverage, or the percentage of the poor that the program reaches.

These limitations can be overcome by a modified version of the approach described above that relies on a household- rather than facility-based survey. A household survey can generate a set of data containing the full range of information needed to produce an equity assessment by collecting two types of information: first, about the household's wealth or assets, using the questions in the left-hand column of the attached questionnaire;²² and second, about the household members' use of or exposure to the services provided by the program(s) of interest. The collected data can be analyzed in either (or both) of two ways, depending on the type of information desired:

- One way would be to use only data from the household survey. The procedure would be analogous to that for a DHS survey employed in this report:
 - Asset information from the survey-generated data set would be used as the basis for the construction of a wealth index, weighting the individual items using some method like principal components analysis.
 - The individuals in the sample would be ranked in order of the index values for their households, then divided into groups like quintiles.
 - The coverage rates in each quintile for the service of interest would be calculated.
- A second approach would be to use the weights for each item appearing on the attached questionnaire in determining the wealth of each individual, instead of calculating the weights from the new household data set. Once the individuals' wealth is determined, the individuals would be ranked, divided into quintiles, and the coverage rate in each quintile would be calculated. In the case of programs undertaken in only one region of the country, it would

²² Or, if one is willing to forego the benefits of the second analytical approach described below, using any of several other asset questionnaires that exist. Examples include the INDEPTH health equity survey tool (available at: www.indepth-network.org/core_documents/indepthtools.htm) or the model questionnaire developed by M. Mahood Khan and David Hotchkiss of the PHR Plus project (which can be found at: www.phrplus.org).

provide a comparison of the economic status of the people served with that of the entire country rather than of only the region where the programs are active;²³ in the case of nationwide as well as regional programs, it would permit a comparison with the other service programs covered in this report.

²³ This additional perspective could be particularly helpful in an assessment of a program seeking to reach the poor by focusing on especially backward districts. A report presenting only a finding that the program was reaching the better-off people in those districts could produce an impression that it had failed to reach its intended beneficiaries. But a comparison between the economic status of the program's beneficiaries with that of the national population might well reveal that most of the beneficiaries were poor by national standards and that the program was thus considerably more successful than otherwise thought.

Mali
2001 - ASSET QUESTIONNAIRE

Question	Score if "Yes"	Score if "No"	Item Score
<i>1. In your household, is/are there?</i>			
Electricity	0.39356	-0.04774	_____
One or more radios	0.02727	-0.05671	_____
One or more televisions	0.32874	-0.04849	_____
One or more refrigerators	0.54804	-0.02946	_____
One or more bicycles	-0.02269	0.01836	_____
One or more motorcycles, scooters	0.09121	-0.02116	_____
One or more cars, trucks	0.44170	-0.01890	_____
One or more telephones	0.63760	-0.01519	_____
One or more horse-drawn carts	-0.08594	0.05520	_____
One or more plows	-0.08791	0.07127	_____
One or more horses	-0.10564	0.01049	_____
One or more wheelbarrows	-0.04371	0.00072	_____
One or more donkeys	-0.08381	0.06552	_____
<i>2. Does your household have a domestic worker not related to head?</i>	0.64585	-0.00758	_____
<i>3. Do the members of your household work their own or family's agricultural land?</i>	-0.08233	0.03124	_____
<i>4. What is the principal source of drinking water for your household?</i>			
Piped water in residence	0.42921	-0.02412	_____
Piped water in the building	0.25645	-0.00605	_____
Piped water in public faucet	0.07237	-0.01639	_____
Open well in residence	-0.01062	0.00109	_____
Open well in the building	-0.03830	0.00146	_____
Public open well	-0.08265	0.04961	_____
Covered well in residence	-0.01475	0.00065	_____
Covered well in the building	0.00392	-0.00007	_____
Public covered well	-0.05873	0.00567	_____
River	-0.03216	0.00236	_____
Lake	-0.07630	0.00049	_____
Canal	-0.14554	0.00001	_____
Springwater	-0.06018	0.00046	_____
Tanker truck	0.32589	-0.00005	_____
Bottled water	0.04757	-0.00004	_____
Other	0.03792	-0.00005	_____
<i>5. What is the principal source of fuel for cooking used by your household?</i>			
Electricity	0.46785	-0.00011	_____
Gas	0.49208	-0.00241	_____
Biogas	0.35100	-0.00006	_____
Kerosene	0.11724	-0.00003	_____
Coal	0.36288	-0.00074	_____
Charcoal	0.28371	-0.03135	_____
Wood	-0.03759	0.17710	_____
Dung, manure	-0.02197	0.00125	_____
Other	0.03935	-0.00051	_____

Mali
2001 - ASSET QUESTIONNAIRE

Question	Score if "Yes"	Score if "No"	Item Score
<i>6. What is the principal type of toilet facility used by your household?</i>			
Private flush toilet in residence	0.23422	-0.00975	_____
Shared flush toilet in residence	0.15938	-0.00407	_____
Private pit latrine	-0.03299	0.01363	_____
Shared pit latrine	0.01562	-0.00619	_____
Ventilated improved pit latrine	0.14296	-0.00583	_____
Shared ventilated improved pit latrine	0.14845	-0.00884	_____
Bush, field as latrine	-0.08440	0.02995	_____
Other	0.03989	-0.00001	_____
<i>7. What is the principal material used for the floors in your household?</i>			
Dirt, sand, dung	-0.05856	0.29978	_____
Cement	0.01107	-0.00002	_____
Wood plank	0.08809	-0.00006	_____
Parquet, polished wood	0.67836	-0.00869	_____
Tile	0.18538	-0.00008	_____
Vinyl, asphalt strips	0.27254	-0.04541	_____
Brick	0.42858	-0.00122	_____
Other	0.03830	-0.00001	_____
Total Household Asset Score (sum of individual item scores)			_____

2001 - QUINTILE CUT-OFF POINTS

Wealth Quintile	Asset Index Value	
	Bottom Cut-Off	Top Cut-Off
Low	Low	-0.64343
Second	-0.64343	-0.42951
Third	-0.42951	-0.18055
Fourth	-0.18055	0.53873
High	0.53873	High

ANNEX C. COUNTRIES COVERED BY THE HNP - POVERTY REPORT PROJECT*

East Asia and Pacific

Cambodia	2000
Indonesia	1997, 2002-03
Philippines	1998, 2003
Vietnam	1997, 2002

Europe and Central Asia

Armenia	2000
Kazakhstan	1995, 1999
Kyrgyz Rep.	1997
Turkey	1993, 1998
Turkmenistan	2000
Uzbekistan	1996

Latin America and the Caribbean

Bolivia	1998, 2003
Brazil	1996
Colombia	1995, 2000, 2005
Dominican Rep.	1996, 2002
Guatemala	1995, 1998-99
Haiti	1994-95, 2000
Nicaragua	1997- 98, 2001
Paraguay	1990
Peru	1996, 2000

Middle East and North Africa

Egypt	1995, 2000
Jordan	1997
Morocco	1992, 2003-04
Yemen	1997

South Asia

Bangladesh	1996-97, 1999-2000, 2004
India	1992-93, 1998-99
Nepal	1996, 2001
Pakistan	1990-91

Sub-Saharan Africa

Benin	1996, 2001
Burkina Faso	1992-3, 1998-9, 2003
Cameroon	1991, 1998, 2004
Central African Rep.	1994-95
Chad	1996-97, 2004
Comoros	1996
Cote d'Ivoire	1994
Eritrea	1995
Ethiopia	2000
Gabon	2000
Ghana	1993, 1998, 2003
Guinea	1999
Kenya	1993, 1998, 2003
Madagascar	1997
Malawi	1992, 2000
Mali	1995-96, 2001
Mauritania	2000-01
Mozambique	1997, 2003
Namibia	1992, 2000
Niger	1998
Nigeria	1990, 2003
Rwanda	2000
Senegal	1997
South Africa	1998
Tanzania	1996, 1999, 2004
Togo	1998
Uganda	1995, 2000-01
Zambia	1996, 2001-02
Zimbabwe	1994, 1999

* Note: electronic versions of reports for all countries are currently available at: www.worldbank.org/povertyandhealth/countrydata. While supplies last, paper copies may be obtained at no charge by sending a request to the World Bank's health advisory service: healthpop@worldbank.org.